

Publication status: Not informed by the submitting author

# Female homicides in the state of São Paulo, Brazil: a time series analysis from 1980 to 2022

Edson Zangiacomi Martinez, Claudia Benedita dos Santos, Elisangela Ap. da Silva Lizzi, Miriane Lucindo Zucoloto

<https://doi.org/10.1590/SciELOPreprints.12427>

Submitted on: 2025-06-28

Posted on: 2026-04-20 (version 2)  
(YYYY-MM-DD)

Version justification: Versão após pareceres da avaliação por pares.

**Female homicides in the state of São Paulo, Brazil: a time series analysis from 1980 to 2022**

**Homicídios femininos no estado de São Paulo, Brasil: uma análise de série temporal de 1980 a 2022**

**Short title:** Female homicides in the state of São Paulo, Brazil

Edson Zangiacomi Martinez<sup>I</sup>, Cláudia Benedita dos Santos<sup>II</sup>, Elisângela Ap. da Silva Lizzi<sup>III</sup>, Miriane Lucindo Zucoloto<sup>IV</sup>

I.Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Ribeirão Preto, SP, Brazil. E-mail: [edson@fmrp.usp.br](mailto:edson@fmrp.usp.br). ORCID: <https://orcid.org/0000-0002-0949-3222>

II.Universidade de São Paulo, Escola de Enfermagem de Ribeirão Preto, Ribeirão Preto, SP, Brazil. E-mail: [cbsantos@eerp.usp.br](mailto:cbsantos@eerp.usp.br). ORCID: <https://orcid.org/0000-0001-7241-7508>

III.Universidade Tecnológica Federal do Paraná, Cornélio Procópio, PR, Brazil. E-mail: [elisangelalizzi@gmail.com](mailto:elisangelalizzi@gmail.com). ORCID: <https://orcid.org/0000-0001-7064-263X>

IV.Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Ribeirão Preto, SP, Brazil. E-mail: [mirianezucoloto@gmail.com](mailto:mirianezucoloto@gmail.com). ORCID: <https://orcid.org/0000-0002-4745-227X>

**Corresponding author:** Edson Zangiacomi Martinez, Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Av. Bandeirantes 3900, 14049-900, Ribeirão Preto (SP), Brazil, [edson@fmrp.usp.br](mailto:edson@fmrp.usp.br).

**Conflict of interests:** nothing to declare.

**Funding:** CNPq, grant number 305206/2023-0.

**Approval by Research Ethics Committee:** This study analyses aggregated data from a public health information system and does not identify individuals. Therefore, approval by a Research Ethics Committee is not required.

**Data Availability Statement:** The data used in this article can be found in open-access health information systems that are free to use.

**Authors' contributions:** EZM: Conceptualization, Data curation, Project administration, Supervision, Software, Formal analysis, Visualization, Writing –original draft. CBS: Visualization, Writing –review & editing. EAS: Methodology, Software, Formal analysis, Writing –review & editing. MLZ: Methodology, Visualization, Writing –review & editing.

## ABSTRACT

**Objective:** This ecological study analyzes the temporal evolution of female homicide rates in the state of São Paulo, Brazil, from 1980 to 2022, using data from the Mortality Information System (SIM). **Methods:** Possible change points in the trend of these rates were identified using the Bayesian Estimator of Abrupt Change, Seasonality and Trend (BEAST) model. **Results:** A total of 29,234 female homicide cases were recorded during the period, including 3,914 among young women aged 15 to 19. The analysis detected two major change points in 1995 and 2004, dividing the time series into three segments: an increasing trend until 1994, a stable period from 1995 to 2003, and a decreasing trend from 2004 onward. Among young women, the data identified three change points, with a significant and consistent decline in homicide rates from 2004 to 2022. **Conclusion:** The number of female homicides in São Paulo has declined significantly since 2004,

particularly among girls aged 15–19. Although this is linked to policy changes, the decline is likely due to multiple factors. The study highlights the importance of ongoing gender-focused policies and robust analytical tools in guiding public safety efforts.

**Keywords:** Violence against women; Gender-based violence; Brazil; Homicide; Public health surveillance

## RESUMO

**Objetivo:** Este estudo ecológico analisa a evolução temporal das taxas de homicídio feminino no estado de São Paulo, Brasil, de 1980 a 2022, usando dados do Sistema de Informações sobre Mortalidade (SIM). **Métodos:** Possíveis pontos de mudança na tendência dessas taxas foram identificados usando o modelo Bayesian Estimator of Abrupt Change, Seasonality and Trend (BEAST). **Resultados:** Um total de 29.234 casos de homicídio feminino foi registrado durante o período, incluindo 3.914 entre mulheres jovens de 15 a 19 anos. A análise detectou dois pontos de mudança importantes em 1995 e 2004, dividindo a série temporal em três segmentos: uma tendência crescente até 1994, um período estável de 1995 a 2003 e uma tendência decrescente de 2004 em diante. Entre as mulheres jovens, os dados identificaram três pontos de mudança, com um declínio significativo e consistente nas taxas de homicídio de 2004 a 2022. **Conclusões:** O número de homicídios femininos em São Paulo diminuiu significativamente desde 2004, principalmente entre as jovens de 15 a 19 anos. Embora isso esteja ligado a mudanças de políticas, o declínio provavelmente se deve a vários fatores. O estudo destaca a importância de políticas contínuas com foco em gênero e ferramentas analíticas robustas para orientar os esforços de segurança pública.

**Palavras-chave:** Violência contra a mulher; Violência de gênero; Brasil; Homicídio; Vigilância em saúde pública

## Introduction

As a World Health Organization (WHO) report has observed, violence against women is not a new phenomenon, nor are the consequences for their physical, mental, and reproductive health<sup>1</sup>. However, the growing recognition that these acts violate women's and girls' rights, limiting their participation in society and damaging their health and well-being, is a more recent development<sup>1</sup>. This WHO report defines 'violence against women' as encompassing various forms of violence, such as intimate partner violence, rape, sexual assault and other forms of sexual violence perpetrated by individuals other than partners. It also covers female genital mutilation, honor killings and the trafficking of women<sup>1</sup>.

Brazil has implemented various initiatives to combat violence against women. In 2006, Law 11.340/2006 was introduced, also known as the Maria da Penha Law<sup>2</sup>, recognizing gender-based violence as a human rights violation rooted in inequality. Prior to this law, situations of violence against women were, in most cases, considered crimes of lesser offensive potential. The penalties for these crimes were often symbolic, such as the distribution of basic food baskets or community labor, which contributed to a feeling of impunity<sup>3</sup>. In 2015, the Brazilian government formally recognized 'femicide' as a specific and aggravated form of homicide by including this term in Law 13.104/2015 within the penal code, thus categorizing it as a heinous crime<sup>4</sup>.

Despite legislative advances, Brazil remains one of the most violent countries for women<sup>5</sup>. A study using data from the 2019 National Health Survey showed that 19.38% of Brazilian women reported experiencing violence<sup>6</sup>. Psychological violence was the most common subtype, both in isolation and in conjunction with other subtypes. Furthermore, the Global Burden of Disease Study (GBD) on interpersonal violence against women showed that Brazil's homicide rates among women aged 15 to 49 have

remained consistently high between 1990 and 2019, indicating an inability of conservative Brazilian society to protect women<sup>7</sup>.

This article uses time series analysis to examine annual female homicide rates in São Paulo from 1980 to 2022, based on data from the Brazilian Mortality Information System (SIM, in Portuguese). For the purposes of this article, 'female homicide' refers to the murder of women in the broadest sense<sup>8</sup>. This includes cases of femicide, which are related to gender issues, as well as cases involving general delinquency. As the coding of mortality data in the SIM follows the International Classification of Diseases and is based on death certificates which do not specify the circumstances of the crime, analyzing data on female homicides can provide insight into how femicide cases have evolved within the Brazilian population.

## **Methods**

### ***Study design and data source***

This ecological and descriptive study is based on data from the Brazilian Mortality Information System. It considers the annual number of female homicides in São Paulo, Brazil, from 1980 to 2022, categorizing deaths according to place of residence. This period was chosen based on the availability of consolidated data and the fact that it is long enough to make inferences. São Paulo is Brazil's most populous and developed state, accounting for around 22% of the country's population. The study used place of residence as a basis for analysis, since it more accurately reflects the social and cultural environment in which women live. This includes structural inequities and the availability of local protection and support systems, all of which can affect their susceptibility to violence.

In view of the potential demographic changes resulting from a reduction in the proportion of young people in the population over the period, the analyses considered homicides in all age groups, as well as among those in the final phase of adolescence (15-19-year-olds).

Although this age group does not account for the largest proportion of female homicide victims in Brazil<sup>5</sup>, it is characterized by a high incidence of reported interpersonal and self-directed violence. In 2002, the rate of interpersonal and self-directed violence was reported as 707.1 cases per 100,000 inhabitants for women aged 15-19, compared to 585.8 and 503.0 cases for the 20-24 and 25-29 age groups, respectively<sup>9</sup>.

Up to 1995, female homicides were identified using codes from the 9th Revision of the International Classification of Diseases (ICD-9): E960–E978 (homicides and intentional injuries to other people) and E990–E999 (legal interventions). After 1995, the following ICD-10 codes are considered: X85–Y09 (aggression) and Y35 (legal intervention). To calculate the mortality rates, population projections for each federative unit by sex for each year, as well as population counts, provided by the Brazilian Institute of Geography and Statistics (IBGE), were used as the denominator.

Brazil is the fifth largest country in the world by area, spanning a territory of 8,514,876 square kilometers. São Paulo, located in the southeast region, is the most populous state in Brazil, accounting for around a fifth of the country's population. As this study used secondary data and no individuals were identified, it was not submitted to a research ethics committee. This is in accordance with national and international legislation regulating research involving humans.

### ***Statistical analysis***

The Change Point Analysis (CPA) method includes statistical tools used to investigate potential changes occurring within a series of data points. The Bayesian Estimator of Abrupt Change, Seasonal Change and Trend (BEAST) is a flexible CPA method that detects abrupt changes (i.e., change points), cyclic variations, and nonlinear trends in time series observations<sup>10</sup>. Consider a time series denoted by  $(t_i, y_i)$ , where  $i = 1, \dots, n$ , and  $y_i$  are observations of a variable at the time  $t_i$ . In a general manner, BEAST assumes that

this time series comprises three components, trend ( $T$ ), seasonality ( $S$ ), change point and noise. The global model is thus given by

$$y(t_i, \theta) = T(t_i, \theta_T) + S(t_i, \theta_S) + e_i, i = 1, \dots, n,$$

where the parameters  $\theta_T$  and  $\theta_S$  specify the trend and seasonal factors and implicitly encode the abrupt change points. In addition,  $e_i$  are the residuals of the model. By “abrupt”, the authors of the BEAST algorithm refer to “any turning points or breakpoints at which trend or seasonal signals start to deviate from the previous regular trajectories”<sup>10</sup>. The method thus calculates the probability of a change point occurring at each year of the study period, denoted by  $cpPr$ , and the probability distribution of the number of change points in the trend component. The mode of this distribution corresponds to the optimal number of change points. Further details can be obtained from the original article by Zhao et al.<sup>10</sup>.

In the present study, we do not consider the presence of seasonal factors. For estimating the unknown parameters, the Bayesian approach is considered, where the Markov Chain Monte Carlo (MCMC) sampling technique is employed to obtain random samples for posterior inference. Three MCMC chains were simulated, each with 100,000 samples. The first 1,000 samples from each chain were discarded to avoid the effect of the initial values. This analysis has been performed on R (version 4.4.1) using the “Rbeast” package. In the time series intervals in which the BEAST model suggests an approximately linear trend, it is possible to calculate the annual percentage rate change (APC)<sup>11</sup>. This is calculated using the equation  $APC = (e^b - 1)100\%$ , where  $e$  is the base of the natural logarithm and  $b$  is the angular coefficient of the Prais-Winsten regression model. This analysis used the “prais” package in R.

## Results

From 1980 to 2022, a total of 29.234 cases of female homicide were reported in the state of São Paulo, of which 3,914 were among women aged 15 to 19. No female deaths due to legal interventions were recorded until 1995 (ICD-9 codes E990–E999). After that year, 22 deaths (ICD-10 code Y35) were recorded. This indicates that the majority of deaths were caused by homicides and violence.


Figure 1 (panel A) shows the results of using the BEAST algorithm to perform sequence decomposition and change-point analysis on the time series of female homicide rates in the state of São Paulo, where is founded a 69.1% probability that the trend component has two change points. The results indicate two abrupt change points occurring in 2004 and 1995, with respective probabilities (cpPr) of 0.9405 and 0.7742. These two change points divide the trend factors within the time series into three segments. The first segment shows an approximately linear increase from 1980 to 1994, with rates ranging from 2.78 to 4.69 homicides per 100,000 inhabitants and an annual percentage change (APC) of 3.14% (95% confidence interval (CI): 1.72% to 4.59%). The second segment shows an increase from 1995 to 1999, when the homicide rate peaked at 6.18 per 100,000 inhabitants, followed by a decrease until 2003. In a visual interpretation of the graph in Figure 1, if a horizontal line segment is fully included within the 95% credibility interval represented by the shaded area, as occurred between 1995 and 2003, then there is no statistically significant upward or downward trend during that period (similar to  $p < 0.05$ ). Finally, the third segment shows a linear decrease from 2004 until the end of the series. In 2004, the rate of female homicides was 4.25 per 100,000 inhabitants, and by 2022, this rate had fallen to 1.53, the lowest rate recorded during the entire study period (APC - 5.3%, 95%CI: -6.2% to -4.4%).

Figure 1 (panel B) shows the results of applying the BEAST algorithm to the time series of homicide rates among young females (aged 15 to 19) in the state of São Paulo. It was found that there is a 53.3% probability of three change points in the trend component: in 1984 (cpPr 0.2509), 1995 (cpPr 0.8843) and 2004 (cpPr 0.9981). Between 1984 and 1994, the rate of homicides per 100,000 people increased from 6.76 to 7.89, with a not significant APC (1.24%, 95%CI: -0.59% to 3.11%,  $p=0.17$ ). Between 1995 and 2023, the rate varied relatively little. In 1995, the rate was 9.61 homicides per 100,000 people. An observable peak of 10.84 homicides per 100,000 inhabitants was reached in 1997, followed by a decrease until 2003, when the rate equaled 9.70 homicides per 100,000 inhabitants. However, no significant APC was observed between 1995 and 2023 (0.07%, 95% CI: -1.79% to 1.96%,  $p = 0.95$ ), indicating that the rates remained relatively stable during this period. From 2004 to 2022, there was a reduction from 7.42 to 1.93 homicides per 100,000 inhabitants, with an APC of -6.7% (95%CI: -7.53% to -5.86%).

| Period                          | APC (%) | 95%CI            | <i>p</i> value | Homicides per 100,000 inhabitants at the start and finish of the period |
|---------------------------------|---------|------------------|----------------|---|
| <b>All female age groups</b>    |         |                  |                |   |
| 1980 – 1994                     | 3.14    | (1.72 to 4.59)   | <0.01          | 2.78 – 4.69   |
| 1995 – 2003                     | -1.03   | (-3.29 to 1.27)  | 0.33           | 5.63 – 5.14   |
| 2004 – 2022                     | -5.3    | (-6.2 to -4.4)   | <0.01          | 4.25 – 1.53   |
| <b>Young women (aged 15–19)</b> |         |                  |                |   |
| 1980 – 1983                     | 1.38    | (1.37 to 1.39)   | <0.01          | 4.47 – 6.36   |
| 1984 – 1994                     | 1.24    | (-0.59 to 3.11)  | 0.17           | 6.76 – 7.89   |
| 1995 – 2003                     | 0.07    | (-1.79 to 1.96)  | 0.95           | 9.61 – 9.70   |
| 2004 – 2022                     | -6.7    | (-7.53 to -5.86) | <0.01          | 7.42 – 1.93   |

**Table 1.** Annual percentage change (APC) for each time interval defined in the BEAST analysis.

These results are summarized in Table 1. Details of the results of the BEAST analysis and R codes are provided in the Supplementary Material.



**Figure 1.** The red line shows the observed annual homicide rate for (A) all female age groups and (B) young women (aged 15–19) per 100,000 inhabitants in the Brazilian state of São Paulo from 1980 to 2022. The black line shows the trend estimated using the BEAST algorithm, and the shaded area represents a 95% credibility interval. The dashed vertical lines indicate the change points and cpPr denotes the corresponding probability of occurrence. APC denotes the annual percentage rate change.

## Discussion

Our analyses highlight a significant decrease in female homicide rates in São Paulo, particularly since 2004 and among young women. While other researchers have also observed this reduction<sup>12-18</sup>, the present article provides new insights. First, it

demonstrates the practicality of the BEAST method for trend analysis using accessible software like R. Second, it updates earlier findings with an extended time series (1980–2022), in line with the current data availability in official systems. A third contribution is its focus on female homicides. Although the findings do not specifically address femicide this focus enables a more targeted discussion grounded in gender-specific dynamics and legislation, while considering vulnerabilities often overlooked in analyses of the general population.

Analyzing public safety indicators in São Paulo municipality from 1996 to 2008, Peres et al.<sup>12</sup> propose three main explanatory hypotheses for the homicide reduction: investment in public safety, socio-economic changes with improved quality of life and demographic changes with a reduction in the proportion of young people in the population. Supporting the public safety investment hypothesis, Goertzel and Kahn<sup>13</sup> suggest that the figures in São Paulo city and state between 2001 and 2007 may be attributable to more effective policing, such as the stricter enforcement of gun control.

The authors also noted that, in October 2003, the federal government introduced legislation restricting firearm imports, prohibiting unregistered gun ownership and public carrying, while also increasing penalties for violations of gun control laws. This legislation also established a weapons buy-back program involving federal and municipal governments and civil society<sup>19</sup>. Between July 2004 and October 2005, 470,000 guns were voluntarily surrendered in Brazil for up to \$100 each. At the time, this was considered the second-largest buy-back campaign ever carried out worldwide<sup>19</sup>. Additionally, Peres et al.<sup>14</sup> report a 169.5 per cent increase in public security investment in São Paulo state between 1997 and 2008, with a 219.3% rise from 1997 to 2001 alone. Nadanovsky<sup>15</sup>, meanwhile, attributed the decrease in homicide rates in São Paulo state to

an increase in incarceration, noting that the number of prisoners per 100,000 inhabitants had risen from 182 in 1996 to 341 in 2005.

In addition to the possible impact of changes in the demographic composition of the population on the reduction in homicide mortality rates in São Paulo state, the following factors, which are not related to public safety, have been identified: demographic changes, particularly a decrease in the proportion of young people; investment in social policies with subsequent socioeconomic improvements and enhanced quality of life; greater social participation through consolidated civil society initiatives; a declining unemployment rate; and state investment in education and culture<sup>14,16-18</sup>. Another possible explanation is heightened control by organized criminal groups over local crime<sup>20-22</sup>. The consolidation of a faction's hegemony over the criminal underworld would have helped to establish peace in the markets for drugs, arms, vehicles and contraband, as well as the legal markets associated with them<sup>20</sup>. This would have reduced conflicts and, consequently, the number of homicides.

All cited articles help to explain the general homicide decline from 2004. However, except for Martins et al.<sup>18</sup>, the ecological studies<sup>12-17</sup> focused on the São Paulo's general population, rather than specifically on female homicides. In this specific context, some key events aligning with the beginning of the decline in homicide rates in 2004 include the launch of the National Plan for Policies for Women, stemming from the First National Conference on Policies for Women, which prioritized coordinating and expanding support networks<sup>23</sup>. Also in 2004, a consortium of feminist organizations and experts proposed domestic and family violence legislation, emphasizing the need for comprehensive laws and public policies. These efforts resulted in the Maria da Penha Law, sanctioned in 2006. This law marks a turning point in Brazil's efforts to protect

women, introducing tools to prevent abuse, ensuring stronger punishment for offenders, and offering support and protection to women in vulnerable situations<sup>24</sup>.

Nearly two decades after enactment, the Maria da Penha Law remains central to protecting women in Brazil. Our analysis shows its 2006 enactment did not cause an abrupt shift in São Paulo's female homicide trend. However, significant national mobilization in support of the law had begun by 2004, and the subsequent updates, including provisions for immediate arrest, preventive detention, tougher sentences, and protective measures, have likely reinforced the steady decline in rates observed since that year.

In addition to these legal protections, in March 2015, the Femicide Law (13,104/2015) was enacted, classifying femicide as an aggravated form of homicide defined as the killing of a woman due to her gender. The law specifically recognizes cases involving domestic and family violence, or acts of contempt or discrimination based on the victim's gender.<sup>24</sup> However, similar to the Maria da Penha Law, the Femicide Law did not result in an immediate shift in São Paulo's female homicide trend in our analysis. While legal protections may affect mortality trends and other outcomes, this change is not expected to be immediate. This is particularly true given the limitations of the public system in enforcing the laws and ensuring citizen safety. Thus, using time series to assess the impact of laws on mortality rates can be challenging, although the enactment of the Femicide Law is undeniably important in classifying cases of domestic and family violence or contempt and discrimination against the female condition as a criminal conduct, thereby sustaining the gradual decline in homicide rates within the state.

The identification of a changepoint in the time series in 1997 coincides with the period immediately following the implementation of ICD-10 in Brazil. In the case of SIM, the transition from ICD-9 to ICD-10 was not merely a change in codes, but an institutional

effort to improve data quality, which reduced the use of ill-defined causes. Additionally, this change point coincides with the peak of the 'homicide epidemic' in the state of São Paulo in the late 1990s<sup>25</sup>. This period was characterized by intense urban violence and the consolidation of organized crime, which disproportionately affected young people. The high rates among women aged 15–19 in 1997 also highlight the absence of specific protective legislation at the time. The Maria da Penha Law, which aimed to address domestic and gender-based violence, and the Disarmament Statute, were only enacted in 2006 and 2003 respectively<sup>2</sup>.

Some authors have discussed the possibility that the Covid-19 pandemic may have contributed to an increase in violent crimes against women in various contexts and geographical regions<sup>26,27</sup>. According to Marques et al.<sup>28</sup>, factors linked to social isolation that contributed to the increase in these crimes include the victim's isolation, making her more vulnerable, the aggressor's use of alcohol or illicit drugs, which intensifies violence, greater ease for the aggressor to control the victim, and unemployment. A study by Nunes et al.<sup>29</sup> showed that the mortality rate due to intentional violent crimes against women in Alagoas, Brazil, was higher in 2020 (5.33 per 100,000 people) than in the years prior to the pandemic. Other Brazilian studies include that by Leite et al.<sup>30</sup>, which showed that psychological, physical, and sexual violence perpetrated by the intimate partners during the pandemic was widespread among women living in Vitória, in the Brazilian state of Espírito Santo. However, this trend does not appear in our data.

This study has several potential limitations. Firstly, and perhaps most importantly, the mortality rates found in this study may be biased due to underreporting issues, as well as the fact that they are based on population projections provided by the IBGE for each year, which may differ from actual census data. Another limitation lies in the fact that we were unable to ascertain the type of homicide, the circumstances surrounding it, or the weapon

used. For example, analyzing homicides by type of weapon can reveal the potential effect of firearm possession legislation on violent deaths. Cerqueira and Lins<sup>31</sup> highlight the possibility of there being a number of 'hidden homicides' within the mortality information system, i.e. homicides that are not officially recorded as such. To address this issue, the authors propose using machine learning techniques to estimate the probability that a violent death from an undetermined cause was actually a homicide. However, this strategy was not employed in the present study. Furthermore, when considering the number of unrecorded homicides in the country, it is necessary to consider cases where there is no death certificate, for example when perpetrators conceal their victims' bodies. This could also introduce bias into the results of this study, particularly given that the occurrence of this type of homicide may have increased over the period considered.

Despite these limitations, our study also presents some strengths. The temporal coverage of over four decades (1980–2022) enables the identification of long-term trends and structural shifts in female homicide rates, and it includes recent data not typically available in earlier studies, extending the analysis beyond previous research, which often ends in the early 2000s or 2010s<sup>12-18</sup>. Using the BEAST model is a methodological strength because, compared to more conventional linear models (e.g. joinpoint regression), BEAST can detect nonlinear trends and multiple change points. It also allows for a probabilistic interpretation of changes and accommodates model uncertainty in trend estimation. The results obtained can inform evidence-based public policy regarding violence against women.

In conclusion, this study describes a substantial and enduring decrease in the rate of female homicides in the Brazilian state of São Paulo, particularly since 2004, with a more pronounced reduction observed among adolescent females aged 15 to 19. Although the observed trends coincide with major legislative and policy developments, female

homicide is a complex phenomenon influenced by multiple interrelated factors. These factors may include demographic changes, socio-economic improvements, shifts in criminal dynamics, civil society engagement and evolving gender norms. While the statistical analysis provides evidence of significant temporal shifts, it does not allow for causal inferences. Nevertheless, the study highlights the importance of continued investment in gender-sensitive policies and emphasizes the value of advanced analytical tools for monitoring public health and safety indicators in complex social contexts.

## References

1. World Health Organization. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. [Internet] Geneva: World Health Organization; 2013. [cited 2025 June 1]. Available from: <https://www.who.int/publications/i/item/9789241564625>
2. Lira MGC, Dos Santos MCB, Cruz MB, de Lacerda Filho EC, de Souza CAC, Paiva FJL, de Almeida JAT. Brazilian government strategies to protect women. *Behav Soc Issues* 2021;30(1):446-64. <https://doi.org/10.1007/s42822-021-00060-4>
3. Meneghel SN, Mueller B, Collaziol ME, de Quadros MM. Repercussions of the Maria da Penha law in tackling gender violence. *Cien Saude Colet* 2013;18(3):691-700. <https://doi.org/10.1590/S1413-81232013000300015>
4. Souza LA, Barros PP. Controversial issues regarding femicide law (Law no. 13.104/2015). *R Fac Dir Univ São Paulo* 2016;111: 263-279.
5. Meira KC, Simões TC, Guimarães RM, Beserra da Silva PG, Mendonça AB, Cristina de Jesus J, Covre-Sussai M. Female homicides in Brazil and its major regions

- (1980-2019): An analysis of age, period, and cohort effects. *Violence Against Women* 2024;30(15-16):3917-42. <https://doi.org/10.1177/10778012231183657>
6. Vasconcelos NM, Gomes CS, Souza JB, Andrade FMD, Bernal RTI, Machado EL, Ribeiro AP, Malta DC. Who are the adult women exposed to violence in Brazil? *Rev Saude Publica* 2025;59:e8. <https://doi.org/10.11606/s1518-8787.2025059005701>
  7. Pinto IV, Vasconcelos NM, Corassa RB, Naghavi M, Marinho F, Malta DC. Mortality and years of life lost to death or disability by interpersonal violence against women in Brazil: Global Burden of Disease Study, 1990 and 2019. *Rev Soc Bras Med Trop* 2022;55(suppl 1):e0287. <https://doi.org/10.1590/0037-8682-0287-2021>
  8. Caicedo-Roa M, Bandeira LM, Cordeiro RC. Femicide and Feminicide discussing and expanding the concept. *Rev Estud Fem* 2022;30(3):e83829. <https://doi.org/10.1590/1806-9584-2022v30n383829>
  9. Fundação Oswaldo Cruz. Ciclo de informes sobre a situação de saúde da juventude brasileira. Informe I: violências e acidentes. [Internet] Rio de Janeiro: Fundação Oswaldo Cruz; 2025. [cited 2026 Jan 20]. Available from: <https://fiocruz.br/documento/2025/08/informe-i-violencias-e-acidentes-ciclo-de-informes-sobre-situacao-de-saude-da>
  10. Zhao K, Wulder MA, Hu T, Bright R, Wu Q, Qin H, Li Y, Toman E, Mallick B, Zhang X, Brown M. Detecting change-point, trend, and seasonality in satellite time series data to track abrupt changes and nonlinear dynamics: A Bayesian ensemble algorithm. *Remote Sens. Environ* 2019;232:111181. <https://doi.org/10.1016/j.rse.2019.04.034>

11. Clegg LX, Hankey BF, Tiwari R, Feuer EJ, Edwards BK. Estimating average annual per cent change in trend analysis. *Stat Med* 2009;28(29):3670-82. <https://doi.org/10.1002/sim.3733>
12. Peres MF, de Almeida JF, Vicentin D, Ruotti C, Nery MB, Cerda M, Cardia N, Adorno S. Homicide and public security indicator trends in the city of São Paulo between 1996 and 2008: a time-series ecological study. *Cien Saude Colet* 2012;17(12):3249-57. <https://doi.org/10.1590/s1413-81232012001200010>
13. Goertzel T, Kahn T. The great São Paulo homicide drop. *Homicide Stud* 2009;13(4):398-410. <https://doi.org/10.1177/1088767909348711>
14. Peres MF, Vicentin D, Nery MB, de Lima RS, de Souza ER, Cerda M, Cardia N, Adorno S. Decline in homicide rates in São Paulo, Brasil: a descriptive analysis. *Rev Panam Salud Publica*. 2011;29(1):17-26. <https://doi.org/10.1590/s1020-49892011000100003>
15. Nadanovsky P. Increased incarceration rate and reduction in homicides in São Paulo, Brazil, from 1996 to 2005. *Cad Saude Publica* 2009;25(8):1859-64. <https://doi.org/10.1590/s0102-311x2009000800022>
16. Peres MF, de Almeida JF, Vicentin D, Cerda M, Cardia N, Adorno S. Fall in homicides in the city of São Paulo: an exploratory analysis of possible determinants. *Rev Bras Epidemiol* 2011;14(4):709-21. <https://doi.org/10.1590/s1415-790x2011000400017>
17. Gawryszewski VP, Costa LS. Social inequality and homicide rates in Sao Paulo City, Brazil. *Rev Saude Publica* 2005;39(2):191-7. <https://doi.org/10.1590/s0034-89102005000200008>
18. Martins LC, Souza MLM, Silva TPRD, Souza HP, Santos FBO, Dumont-Pena É, Matozinhos FP. Timeline trend of homicides of women in the states of the

- Southeast region of Brazil between 2007 and 2019. *Cien Saude Colet* 2023;28(6):1631-42. <https://doi.org/10.1590/1413-81232023286.16082022>
19. Lucas P. Disarming Brazil: Lessons and Challenges. *NACLA Rep Am* 2008;41(2):27-31 <https://doi.org/10.1080/10714839.2008.11725399>
  20. Biderman C, Mello JMP, Lima, RSD, Schneider A. Pax Monopolista and Crime: The Case of the Emergence of the Primeiro Comando da Capital in São Paulo. *J Quant Criminol* 2019; 35:573-605. <https://doi.org/10.1007/s10940-018-9393-x>
  21. Feltran G, Lero C, Cipriani M, Maldonado J, Rodrigues FDJ, Silva LEL, et al. Variações nas taxas de homicídios no Brasil: Uma explicação centrada nos conflitos faccionais. *Dilemas, Rev Estud Conflito Controle Soc* 2022;15(spe4):311-48. <https://doi.org/10.4322/dilemas.v15nesp4.46920>
  22. Moreira MC, Andrade LT. Reduction of Homicides in São Paulo and Medellín: What Does the Literature Say? *Dilemas, Rev Estud Conflito Controle Soc* 2023;16(2):e51667. <https://doi.org/10.4322/dilemas.v16.51667>
  23. Diniz SG, Silveira LP, Mirim LA. Vinte e cinco anos de respostas brasileiras em violência contra as mulheres no Brasil (1980-2005) - alcances e limites. São Paulo: Coletivo Feminista Sexualidade e Saúde; 2006.
  24. Coimbra JC, Ricciardi U, Levy L. Lei Maria da Penha, equipe multidisciplinar e medidas protetivas. *Arq Bras Psicol* 2018, 70(2):158-72.
  25. Cardia N, Adorno S, Poletto FZ. Homicide rates and human rights violations in São Paulo, Brazil: 1990 to 2002. *Health Hum Rights* 2003, 6(2):14-33. <https://doi.org/10.2307/4065428>
  26. Roesch E, Amin A, Gupta J, García-Moreno C. Violence against women during covid-19 pandemic restrictions. *BMJ* 2020;369:m1712. <https://doi.org/10.1136/bmj.m1712>

27. Sánchez OR, Vale DB, Rodrigues L, Surita FG. Violence against women during the COVID-19 pandemic: An integrative review. *Int J Gynaecol Obstet* 2020;151(2):180-7. <https://doi.org/10.1002/ijgo.13365>
28. Marques ES, Moraes CL, Hasselmann MH, Deslandes SF, Reichenheim ME. Violence against women, children, and adolescents during the COVID-19 pandemic: overview, contributing factors, and mitigating measures. *Cad Saude Publica* 2020;36(4):e00074420. <https://doi.org/10.1590/0102-311X00074420>
29. Nunes DN, Figueredo PCR, Vitoriano TA, Santos GKRS, de Albuquerque CML, Matos TS, Matos DUS, do Carmo RF, de Souza CDF. Lethal and intentional violent crimes against women in Alagoas, Northeast Brazil: a comparative study before and during the Covid-19 pandemic. *BMC Womens Health* 2024;24(1):614. <https://doi.org/10.1186/s12905-024-03464-7>
30. Leite FMC, Venturin B, Eduarda Portes Ribeiro L, De Paula Silva R, Luis Alves M, Wehrmeister FC, Santos DF. Intimate partner violence against women during covid-19: A population-based study in Vitória, state of Espírito Santo, Brazil. *PLoS One* 2023;18(12):e0295340. <https://doi.org/10.1371/journal.pone.0295340>
31. Cerqueira DRC, Lins GOA. Mapa dos homicídios ocultos no Brasil entre 1996 e 2021. Brasília, DF: Ipea,2024

## **Supplementary Material to: Female homicides in the state of São Paulo, Brazil: a time series analysis from 1980 to 2022**

This document provides supplementary material to our paper “Female homicides in the state of São Paulo, Brazil: a time series analysis from 1980 to 2022” which was published in the *Revista Brasileira de Epidemiologia* (2026). It contains further details on the methods of analysis and results that are summarized in the main report.

### **The BEAST method**

The BEAST approach is a common method to decompose time series into trend and seasonality and for detecting abrupt changes in these two components. It has been used in a wide range of applications, including studies of land surface temperature<sup>1</sup>, climate change detection<sup>2</sup> and the dynamics of infectious diseases<sup>3</sup>. Compared to joinpoint regression, which is currently widely used in time series studies in epidemiology, the BEAST approach may be more suitable for evaluating the dynamics of some health events, given that in the joinpoint approach, each segment between the change points is linear<sup>4</sup>. In contrast, the BEAST approach can detect nonlinear trends between successive change points using functions such as spline smoothing. In this study, BEAST was shown to be suitable for describing the behavior of female homicide data series, enabling significant periods to be identified.

### **Statistical details**

The Bayesian model averaging time-series decomposition algorithm (BEAST) is a parametric regression method developed by Zhao et al.<sup>5</sup>. It can capture various types of uncertainty, identify abrupt shifts of any size and reveal intricate nonlinear patterns in time series data. Let  $\mathcal{D} = \{t_i, y_i\}_{i=1, \dots, n}$  the observed data, where  $y_i$  denotes observations of a variable at  $n$  time points  $t_i$ . The statistical model for the time series  $y(t_i)$  is given by

$$y(t_i) = S(t_i; \Theta_S) + T(t_i; \Theta_T) + e_i, i = 1, \dots, n,$$

where the parameters  $\Theta_S$  and  $\Theta_T$  are estimated from the data  $\mathcal{D}$ , and they also implicitly encode the abrupt change points. We assume that the noise  $e_i$  is Gaussian with a magnitude of  $\sigma$ , capturing the remainder of the data that is not explained by the seasonal

( $S$ ) and trend ( $T$ ) components<sup>5,6</sup>. To evaluate the unknown parameters, the algorithm employs Bayesian inference.

The R package `Rbeast` can be used to perform this analysis. Consider `wt` a data frame containing the variables `wt$ANO` (years from 1980 to 2022) and `wt$SP` (the female homicide rates in São Paulo from 1980 to 2022).

```
# Data
wt <- data.frame(
  ANO = 1980:2022,
  SP = c(2.78, 3.07, 3.03, 3.94, 4.15, 3.85, 3.76, 4.00, 3.88,
        4.37, 4.34, 4.94, 3.97, 4.39, 4.69, 5.63, 5.54, 5.83,
        5.78, 6.18, 5.98, 5.64, 5.32, 5.14, 4.25, 3.78, 3.80,
        2.84, 3.14, 3.08, 3.13, 2.65, 2.89, 2.78, 2.72, 2.46,
        2.21, 2.14, 1.95, 1.72, 1.67, 1.43, 1.53))
```


The BEAST model can be fitted to data in the following way:

```
# Bayesian changepoint detection algorithm
library(Rbeast)
myts <- ts(wt$SP, start=c(1980), end=c(2022))
out <- beast(myts, season='none',
            dump.ci = TRUE, mcmc.burnin = 1000, mcmc.samples = 100000)
plot(out)
print(out)
```

In this R code, `library(Rbeast)` is used to load the `Rbeast` package, and the `ts` function is used to create the time-series object `myts` containing `wt$SP`. The `beast` function in the `Rbeast` package is then used to fit the BEAST model to the data, where the argument `season='none'` is used to specify that the seasonal component  $S(t_i; \Theta_S)$  will not be used in our analysis, and `dump.ci = TRUE` is used to create credible intervals for the estimated trend component. The argument `mcmc.burnin = 1000` specifies the number of burn-in samples discarded at the start of each chain (the default number of MCMC chains is three), while `mcmc.samples` specifies the number of samples to be collected per MCMC chain.

Figure S1 shows the results obtained from the `beast` function. The dashed vertical lines mark the most likely locations of changepoints and the green curve of  $\text{Pr}(\text{tcp})$  shows the point-wise probability of changepoint occurrence over time. The “`order1`” curve provides an estimate of the mean order of the piecewise polynomials required to adequately fit the trend (the 0-th order is constant and the 1st order is linear). An average order toward 0

means that the trend is more likely to be flat and an order close to 1 means that the trend is linear. “slpSign” are the probabilities of the trend slope being positive (red part), zero (green), or negative (blue). Finally, “error” are the residual errors between the observed and BEAST-modelled values.



**Figure S1.** Results of BEAST of female homicides time series in São Paulo from 2000 to 2022.

The `print(out)` function summarizes and prints the results obtained from the BEAST time series decomposition and segmentation, as shown below.

```

> print(out)

#####
#           Seasonal Changepoints           #
#####
No seasonal/periodic component present (i.e., season='none')

#####
#           Trend Changepoints             #
#####
-----
| Ascii plot of probability distribution for number of chgpts (ncp) |
-----
|Pr(ncp = 0 )=0.000|*
|Pr(ncp = 1 )=0.080|*****
|Pr(ncp = 2 )=0.691|*****
|Pr(ncp = 3 )=0.196|*****
|Pr(ncp = 4 )=0.031|***
|Pr(ncp = 5 )=0.003|*
|Pr(ncp = 6 )=0.000|*
|Pr(ncp = 7 )=0.000|*
|Pr(ncp = 8 )=0.000|*
|Pr(ncp = 9 )=0.000|*
-----
|           Summary for number of Trend ChangePoints (tcp)           |
-----
|ncp_max    = 9    | MaxTrendKnotNum: A parameter you set |
|ncp_mode   = 2    | Pr(ncp= 2)=0.69: There is a 69.1% probability |
|           | that the trend component has 2 changepoint(s). |
|ncp_mean   = 2.19 | Sum{ncp*Pr(ncp)} for ncp = 0,...,9 |
|ncp_pct10  = 2.00 | 10% percentile for number of changepoints |
|ncp_median = 2.00 | 50% percentile: Median number of changepoints |
|ncp_pct90  = 3.00 | 90% percentile for number of changepoints |
-----
| List of probable trend changepoints ranked by probability of |
| occurrence: Please combine the ncp reported above to determine |
| which changepoints below are practically meaningful |
-----
|tcp#          |time (cp)          |prob(cpPr)          |
|-----|-----|-----|
|1             |2004.000000        |0.94046             |
|2             |1995.000000        |0.77420             |
|3             |1992.000000        |0.11254             |
|4             |1984.000000        |0.09895             |
|-----|-----|-----|

```

This output shows that there is a 69.1% probability that the trend component has two change points (the mode for the probability distribution for number of change points, denoted by `ncp_mode`). The locations where these two change points occur and the probabilities associated with each change point are 2004 (94.06%) and 1995 (77.42%).

### Statistical details of APC estimation

When the BEAST model suggests an approximately linear trend in an interval on the time series, it is possible to calculate the correspondent annual percentage rate change (APC)<sup>7</sup>. APC is defined as

$$APC = \frac{\gamma_{j+1} - \gamma_j}{\gamma_j},$$

where  $\gamma_j$  denotes the expected rate at time  $j$ . If we assume that  $\log(\gamma_j)$  is linear over the entire time interval  $[t_0, t_1]$ , then

$$\log \gamma_j = a + bt_j,$$

where  $a$  and  $b$  are constants. Note that

$$\frac{\gamma_{j+1}}{\gamma_j} = \frac{e^{a+bt_{j+1}}}{e^{a+bt_j}} = e^{b(t_{j+1}-t_j)} = e^b$$

when  $t_{j+1} - t_j = 1$ , i.e. when considering a change of one time unit. Therefore,

$$APC = \frac{\gamma_{j+1} - \gamma_j}{\gamma_j} = \frac{\gamma_{j+1}}{\gamma_j} - 1 = e^b - 1,$$

where  $e$  is the base of the natural logarithm and  $b$  is the angular coefficient of the Prais-Winsten regression model.

We let  $y_t$  denote observations of a variable at  $n$  time points  $t = 1, \dots, n$ . A linear regression model is given by

$$\log y_t = a + bt + \epsilon_t,$$

where  $a$  and  $b$  are unknown coefficients and  $\epsilon_t$  is a mean zero error representing other determinants of the outcome. We assume that  $t$  is independent of  $\epsilon_t$ . The Prais-Winsten method involves estimating the correlation between the error at  $t$  and  $t - 1$ , assuming that

$$\epsilon_t = \phi_1 \epsilon_{t-1} + \eta_t,$$

where  $\phi_1$  is an autoregressive (AR) parameter with  $|\phi_1| < 1$  and  $\eta_t$  are independent disturbances with zero mean and variance  $\sigma_\eta^2$ . The constraint  $\phi_1$  ensures that the process is stationary, that is, the covariance between  $\epsilon_t$  and  $\epsilon_{t+h}$  is independent of  $t$ , and therefore

that the variance remains constant over time. In our article, this analysis used the function `prais_winsten` of the `prais` package in R.

For example, to estimate the APC of the of female homicide rates time series between  $t_0 = 1980$  and  $t_1 = 1994$ , we can use the following R code:

```
# APC estimation based on the Prais-Winsten regression model
# 1980 to 1994
library(prais)
pw <- c()
ay <- 1980
by <- 1994
NY <- length(ay:by)
seqt <- (ay-1979):(by-1979)
pw_sample <- data.frame("x" = wt$ANO[seqt], "y" = wt$SP[seqt],
                        "time" = seqt)
pw <- prais_winsten(log(y) ~ x, data = pw_sample, index = "time")
a <- summary(pw)
b1 <- a$coefficients[2,1] # angular coefficient
se <- a$coefficients[2,2] # standard errors
APC <- exp(b1)-1 # APC
L_APC <- exp(b1-se*qt(1-0.05/2,NY-1))-1
U_APC <- exp(b1+se*qt(1-0.05/2,NY-1))-1
message("APC ", round(100*APC,2), " 95%CI (", round(100*L_APC,2),
        " , ", round(100*U_APC,2), ")")
```

In this R code, the variable `wt$ANO` is a vector from the data frame `wt` containing the years 1980 to 2022.

The following output is then obtained for the Prais-Winsten regression model analysis:

```
Call:
prais_winsten(formula = log(y) ~ x, data = pw_sample, index = "time")

Residuals:
    Min       1Q   Median       3Q      Max
-0.133876 -0.057356 -0.002484  0.053558  0.158118

AR(1) coefficient rho after 6 iterations: 0.2012

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) -60.152761  12.923666  -4.654 0.000451 ***
x              0.030957   0.006504   4.760 0.000373 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.09134 on 13 degrees of freedom
```

```
Multiple R-squared: 0.428, Adjusted R-squared: 0.384
F-statistic: 9.727 on 1 and 13 DF, p-value: 0.008146

Durbin-Watson statistic (original): 1.473
Durbin-Watson statistic (transformed): 1.856
```

We can note that the estimated values for the parameters  $\phi_1$  and  $b$  are 0.2012 and 0.030957, respectively. Thus, the APC is estimated to be

$$\text{APC} = (e^{0.030957} - 1)100\% \cong 3.14\%.$$

The lower and upper limits of the 95% confidence interval for the APC can be obtained using the following expressions, respectively:

$$[\exp(b - \text{se}(b) \times t_{(1-0.05/2, \text{df})}) - 1]100\%$$

and

$$[\exp(b + \text{se}(b) \times t_{(1-0.05/2, \text{df})}) - 1]100\%$$

In these expressions,  $\text{se}(b)$  is an estimate of the standard error of  $b$ . In the R output, we have  $\text{se}(b)$  equal to 0.006504. In addition,  $t_{(1-0.05/2, \text{df})}$  is the  $(1-0.05/2)$  percentile of a Student's  $t$  distribution with degrees of freedom ( $\text{df}$ ) given by  $n - 1$ , where  $n$  is the length of the interval. This percentile is obtained using the R function `qt`.

The R code above also provides the following:

```
APC 3.14 95%CI (1.72 , 4.59)
```

Thus, assuming that the annual rate changes are constant throughout the time interval [1980, 1994], we estimate an annual percentage rate change of 3.14% for this period.

R program version 4.1.1 was used for all these analyses.

## References

1. Li J, Li ZL, Wu H, You N. Trend, seasonality, and abrupt change detection method for land surface temperature time-series analysis: Evaluation and improvement. *Remote Sens Environ* 2022; 280, 113222. <https://doi.org/10.1016/j.rse.2022.113222>

2. Dan'azumi S, Mamudu L, Aldrees A. Climate change detection and attribution: Bayesian estimation of abrupt change, seasonality and trend model, and Mann–Kendall trend test approaches. *J Water Clim Chang* 2025;16(5):1895-911. <https://doi.org/10.2166/wcc.2025.004>
3. Yang L, Wang C, Zhou P, Xie N, Tian M, Wang K. Change point detection in brucellosis time series from 2010 to 2023 in Xinjiang China using the BEAST algorithm. *Scientific Reports* 2025;15(1):3830. <https://doi.org/10.1038/s41598-025-88508-0>
4. Kim HJ, Chen HS, Byrne J, Wheeler B, Feuer EJ. Twenty years since Joinpoint 1.0: Two major enhancements, their justification, and impact. *Stat Med* 2022;41(16):3102-30. <https://doi.org/10.1002/sim.9407>
5. Zhao K, Wulder MA, Hu T, Bright R, Wu Q, Qin H, Li Y, Toman E, Mallick B, Zhang X, Brown M. Detecting change-point, trend, and seasonality in satellite time series data to track abrupt changes and nonlinear dynamics: A Bayesian ensemble algorithm. *Remote Sens. Environ* 2019; 232: 111181. <https://doi.org/10.1016/j.rse.2019.04.034>
6. Yang L, Wang C, Zhou P, Xie N, Tian M, Wang K. Change point detection in brucellosis time series from 2010 to 2023 in Xinjiang China using the BEAST algorithm. *Scientific Reports* 2025;15(1):3830. <https://doi.org/10.1038/s41598-025-88508-0>
7. Clegg LX, Hankey BF, Tiwari R, Feuer EJ, Edwards BK. Estimating average annual per cent change in trend analysis. *Stat Med* 2009;28(29):3670-82. <https://doi.org/10.1002/sim.3733>

This preprint was submitted under the following conditions:

- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.