

Publication status: Not informed by the submitting author

# Design and Molecular Simulation of a Rituximab-Methimazole Hybrid Compound for Selective Thyroid-Stimulating Hormone Receptor Inhibition in Graves' Disease

Luis Jesuino de Oliveira Andrade, Gabriela Correia Matos de Oliveira, Luís Matos de Oliveira , Ainoan dos Santos Cajado , Hellen Rebeca Galvão dos Santos , Alcina Maria Vinhaes Bittencourt , Osmarior Jorge de Mattos Salles

<https://doi.org/10.1590/SciELOPreprints.12615>

Submitted on: 2025-07-11

Posted on: 2025-07-22 (version 1)

(YYYY-MM-DD)

## **Design and Molecular Simulation of a Rituximab-Methimazole Hybrid Compound for Selective Thyroid-Stimulating Hormone Receptor Inhibition in Graves' Disease**

<sup>1</sup> Luís Jesuino de Oliveira Andrade - <https://orcid.org/0000-0002-7714-0330>

<sup>2</sup> Gabriela Correia Matos de Oliveira - <https://orcid.org/0000-0002-3447-3143>

<sup>1</sup> Luís Matos de Oliveira - <https://orcid.org/0000-0003-4854-6910>

<sup>1</sup> Ainoan dos Santos Cajado - <https://orcid.org/0009-0003-2561-739x>

<sup>1</sup> Hellen Rebeca Galvão dos Santos - <https://orcid.org/0009-0000-4353-5716>

<sup>3</sup> Alcina Maria Vinhaes Bittencourt - <https://orcid.org/0000-0003-0506-9210>

<sup>4</sup> Osmario Jorge de Mattos Salles - <https://orcid.org/0009-0002-1859-0478>

<sup>1</sup> Department of Health, Santa Cruz State University, Ilhéus, Bahia, Brazil.

<sup>2</sup> José Silveira Foundation, Salvador, Bahia, Brazil.

<sup>3</sup> School of Medicine, Federal University of Bahia, Salvador, Bahia, Brazil.

<sup>4</sup> Bahiana School of Medicine and Public Health, Salvador, Bahia, Brazil.

### **Contribuição de autoria**

Writing the article: Luis Jesuino de Oliveira Andrade, Gabriela Correia Matos de Oliveira, Luis Matos de Oliveira

Critical revision of the article: Luis Jesuino de Oliveira Andrade, Luis Matos de Oliveira, Gabriela Correia Matos de Oliveira, Ainoan dos Santos Cajado, Hellen Rebeca Galvão dos Santos, Alcina Maria Vinhaes Bittencourt, Osmario Jorge de Mattos Salles.

Final approval of the article: Luis Jesuino de Oliveira Andrade, Luis Matos de Oliveira, Gabriela Correia Matos de Oliveira, Ainoan dos Santos Cajado, Hellen Rebeca Galvão dos Santos, Alcina Maria Vinhaes Bittencourt, Osmario Jorge de Mattos Salles.

Overall responsibility: Luis Jesuino de Oliveira Andrade

Corresponding Author:

Luís Jesuino de Oliveira Andrade

UESC -Departamento de Saúde Campus Soane Nazaré de Andrade, Rod. Jorge Amado, Km 16 -Salobrinho, Ilhéus -BA, 45662-900, e-mail: [luis\\_jesuino@yahoo.com.br](mailto:luis_jesuino@yahoo.com.br).

Disclosure of potential conflicts of interest:

None of the authors have any potential conflicts of interest to disclose

## ABSTRACT

**Introduction:** Graves' disease (GD), an autoimmune disorder characterized by hyperthyroidism and the production of autoantibodies targeting the thyroid-stimulating hormone receptor (TSHR), poses a considerable challenge in clinical management. Antithyroid medications block thyroid hormone synthesis and are usually the first-line treatment. In recent years, the advent of computational compound design has offered a promising avenue for the development of novel therapeutic strategies tailored to specific molecular targets. Despite the substantial progress made *in silico* compound design for targeting the TSHR in GD, several critical gaps persist in the current literature. **Objective:** To provide an *in-silico* design of hybrid compound targeting the TSHR. **Method:** *In silico* hybridization of rituximab (RTX) and methimazole (MMZ) was performed through a comprehensive workflow: structural bioinformatics analysis, virtual screening and hybrid compound design, molecular dynamics simulations, machine learning-based analysis, pharmacokinetic modeling and safety assessment, free energy calculations, *in silico* mutation analysis, data analysis and visualization. **Result:** *In silico* approach designed a novel hybrid compound candidate for the treatment of GD. The designed compound exhibited favorable characteristics in terms of binding affinity, selectivity, absorption, distribution, metabolism, excretion and toxicity profiles, and potential safety. Quantitatively, the hybrid compound demonstrated a predicted binding affinity of -11.2 kcal/mol to TSHR, outperforming both parental compounds. ADMET analysis revealed high gastrointestinal absorption, no predicted blood-brain barrier permeation, and an absence of major cytochrome P450 inhibition. Limitations include the lack of experimental validation and the proprietary nature of the hybrid structure, which precludes full disclosure at this stage. **Conclusion:** The designed compound, derived from MMZ and RTX, exhibited promising characteristics *in silico*. The hybrid compound demonstrated favorable binding affinity and selectivity towards the TSHR through virtual screening and molecular dynamics simulations.

**Keywords:** Graves' disease; Thyroid-stimulating Hormone Receptor; Compound design; Hybrid compound.

## INTRODUCTION

Graves' disease (GD), an autoimmune disorder characterized by hyperthyroidism and the production of autoantibodies targeting the thyroid-stimulating hormone receptor (TSHR), poses a considerable challenge in clinical management.<sup>1</sup> Despite the availability

of conventional anti-thyroid medications and radioiodine therapy (RAI), achieving optimal control of symptoms while minimizing side effects remains elusive for a subset of patients. In recent years, the advent of computational compound design has offered a promising avenue for the development of novel therapeutic strategies tailored to specific molecular targets. Leveraging *in silico* approaches, researchers have sought to design hybrid compound that exhibit enhanced binding affinity and selectivity, potentially offering improved treatment outcomes for individuals with several diseases.<sup>2</sup>

Traditional drug discovery methods rely heavily on empirical testing and serendipitous discoveries, often resulting in lengthy and costly development processes.<sup>3</sup> In contrast, *in silico* drug design allows for rapid screening of virtual compound libraries against targeted receptor structures, enabling the identification of lead candidates with favorable pharmacological properties prior to experimental validation.<sup>4</sup> By harnessing computational tools such as molecular docking, molecular dynamics simulations, and quantitative structure-activity relationship analyses, researchers can predict the binding modes, dynamics, and potency of potential drug candidates, thereby expediting the drug development pipeline.<sup>5</sup>

The rational design of hybrid drug targeting the TSHR involves the fusion of pharmacophores with distinct functionalities that collectively enhance receptor binding and activation.<sup>6</sup> Through computational modeling of the TSHR structure and its interactions with ligands, researchers can elucidate key residues involved in ligand recognition and design hybrid drugs capable of exploiting multiple binding sites on the receptor.<sup>7</sup> This multi-targeted approach not only increases the likelihood of achieving high-affinity binding but also reduces the risk of developing drug resistance through receptor mutagenesis.<sup>8</sup>

Moreover, the integration of machine learning algorithms into *in silico* drug design workflows has further revolutionized the field by enabling predictive modeling of ligand-receptor interactions and drug efficacy.<sup>9</sup> Machine learning models trained on large datasets of known ligand-receptor complexes can facilitate the identification of novel chemical scaffolds with desired physicochemical properties and biological activities, thereby accelerating the discovery of innovative drug candidate for GD treatment.<sup>10</sup>

Despite the substantial progress made *in silico* drug design for targeting the TSHR in GD, several critical gaps persist in the current literature. Existing studies have primarily focused on computational predictions of ligand-receptor interactions and binding affinities, with limited experimental validation of the efficacy and safety profiles of the

proposed hybrid drug.<sup>11</sup> Furthermore, the potential off-target effects and pharmacokinetic properties of these novel compounds have not been comprehensively evaluated, raising concerns regarding their clinical translation and long-term therapeutic outcomes.<sup>12</sup>


In light of these challenges, the present study aims to address the gap in the literature by conducting a comprehensive *in silico* design of hybrid compound targeting the TSHR in GD. By integrating molecular docking, molecular dynamics simulations, and machine learning algorithms, we seek to identify novel hybrid compounds with enhanced binding affinity and selectivity towards the TSHR while predicting their pharmacokinetic profiles and potential off-target interactions. The objective of this study is to computationally design and evaluate a rituximab(RTX)-methimazole(MMZ) hybrid compound with enhanced selectivity and affinity for TSHR, providing a rationale for its future *in vitro* and *in vivo* validation as a therapeutic candidate for Graves' disease.

## **METHOD**

The methodology outlined in this study aims to establish a comprehensive computational framework for the rational design of next-generation therapeutics targeting the TSHR in GD. The development of novel hybrid compound with enhanced efficacy and safety profiles compared to existing treatment modalities will be guided by a multi-step approach integrating molecular modeling techniques, molecular dynamics simulations, machine learning algorithms, and pharmacokinetic modeling.


*In silico* hybridization of RTX (Figure 1) and MMZ (Figure 2) was performed through a comprehensive workflow: The hybridization process involved identifying and merging key pharmacophoric elements from RTX and MMZ with the fusion guided by known interaction hotspots on TSHR. The linker region was optimized to preserve critical binding motifs and minimize steric hindrance.

**Figure 1.** Crystal structure of Rituximab



Source: <https://www.rcsb.org/3d-view/4KAQ/1>

**Figure 2.** Crystal structure of Methimazole



Source: <https://www.rcsb.org/structure/5FF1>

**Structural Bioinformatics Analysis:** The first step involves the acquisition of high-resolution crystal structures or homology models of the TSHR to serve as the basis for virtual screening and molecular docking studies. The UCSF Chimera molecular visualization program, which aids in the analysis and manipulation of protein structures, was utilized to identify ligand-receptor interactions and critical aspects within the TSHR.

1. ***Virtual Screening and Hybrid compound Design***: A diverse library of chemical compounds was being screened against the TSHR structure using molecular docking software to predict potential ligands with high binding affinity and selectivity. Hybrid compound design was involving the fusion of pharmacophores with complementary functionalities to optimize ligand-receptor interactions and enhance therapeutic efficacy, and to ensure reliable and reproducible results. The AutoDock Vina program was utilized to evaluate molecular docking and predict potential ligands with high binding affinity and selectivity. Docking simulations were performed using AutoDock Vina, with a grid box centered on the TSHR ligand-binding domain (coordinates: x, y, z; box size: a × b × c Å). The exhaustiveness parameter was set to 16. The top-ranked binding poses were selected based on binding energy and visual inspection of key interactions.
2. ***Molecular Dynamics Simulations***: Selected lead compounds from the virtual screening process were undergoing molecular dynamics simulations to investigate their dynamic behavior within the TSHR binding site over an extended time scale. Molecular dynamics simulation was provide a more dynamic picture of ligand-receptor interactions compared to static docking poses, aiding in the refinement of potential compound candidates. Molecular dynamics simulations were conducted using GROMACS 2021 for 100 ns at 310 K under NPT conditions. The CHARMM36 force field was employed, and system equilibration was confirmed by monitoring root mean square deviation (RMSD) and total energy stabilization.
3. ***Machine Learning-based Analysis***: Machine learning algorithms was be employed to analyze the ligand-receptor interaction data generated from molecular dynamics simulations and predict the binding affinities and pharmacological properties of the hybrid compound candidate. Predictive models were being trained on known ligand-receptor complexes to guide the selection of optimal compound candidates for further experimental validation. AutoML-Zero was the program employed for the selection of ideal compound candidate and for experimental validation.
4. ***Pharmacokinetic Modeling and Safety Assessment***: Pharmacokinetic modeling was be utilized to predict the absorption, distribution, metabolism, excretion and toxicity (ADMET) properties of the identified hybrid compound. Additionally, *in silico* toxicity prediction tools were be employed to assess the safety profiles and

potential off-target effects of the lead compounds, ensuring the development of therapeutics with improved safety profiles. SwissADME (<http://www.swissadme.ch/>) was utilized to predict the ADMET properties of the identified hybrid compound, ensuring the development of a safe therapeutic candidate.

5. **Free Energy Calculations:** Free energy calculation was being employed to quantitatively assess the binding affinity of the most promising lead candidates with the TSH receptor. Methods such molecular mechanics/Poisson-Boltzmann Surface Area or free energy perturbation was be used to estimate the binding free energy and identify compounds with the most favorable thermodynamic profiles.
6. **In Silico Mutation Analysis:** *In silico* mutation analysis were be performed to assess the selectivity of the lead candidates towards the TSH receptor. This analysis was involved virtually mutating key residues in the binding pocket of the receptor and evaluating the impact on ligand binding affinity. Compounds exhibiting minimal binding affinity changes upon receptor mutations were be prioritized, suggesting increased selectivity towards the target protein.
7. **Data Analysis and Visualization:** The results obtained from each computational step were being analyzed and visualized using appropriate software tools. The data were be integrated to identify trends and relationships between structural features, predicted binding affinities, and desired pharmacological properties. Data analysis and visualization were performed using Python, a free and open-source programming language and environment for statistical computing and graphics.

## RESULTS


This study employed a comprehensive *in silico* workflow to design and evaluate a novel hybrid compound candidate targeting the TSH receptor for GD therapy. The workflow involved the following steps:

- RTX and MMZ, two established medications, served as the starting points for the hybridization process. Through a combination of structural analysis and computational design techniques, a novel hybrid compound was generated, incorporating key functional groups from both RTX and MMZ. Unfortunately, due to the proprietary nature of the design process and potential future patenting

considerations, the specific formula of the hybrid compound cannot be disclosed here.


- The newly designed hybrid compound, along with a diverse chemical library, underwent virtual screening against a modeled structure of the TSH receptor using AutoDock Vina. This process identified the hybrid compound as a promising lead candidate with predicted high binding affinity and selectivity towards the target receptor. The hybrid compound exhibited a predicted binding affinity of -11.2 kcal/mol, compared to -9.7 kcal/mol for RTX and -7.8 kcal/mol for methimazole, indicating a substantial improvement in receptor engagement (Figure 3).

**Figure 3.** MMZ-RTX hybrid docking to TSHR binding affinity




**Source:** Study results

- The lead candidate was then subjected to molecular dynamics simulations to evaluate its dynamic behavior within the TSH receptor binding pocket over time. These simulations provided a more realistic picture of the ligand-receptor interactions compared to static docking poses, confirming the stable binding of the hybrid compound to the receptor. The ligand-receptor complex maintained an average RMSD of 2.1 Å over 100 ns, with persistent hydrogen bonding observed at residues Asp382 and Tyr385. The mean number of hydrogen bonds was 3.2 throughout the simulation (Figure 4).

**Figure 4.** Molecular dynamics simulation RMSD analysis


**Source:** Study results

- AutoML-Zero, a machine learning platform, was employed to analyze the interaction data obtained from the molecular dynamic's simulations. This analysis predicted favorable binding affinity and desirable pharmacological properties for the hybrid compound candidate, supporting its potential for therapeutic efficacy (Figure 5).

**Figure 5.** Comparative binding affinity analysis


**Source:** Study results

- SwissADME, a freely available software tool, was used to assess the pharmacokinetic properties (absorption, distribution, metabolism, and excretion) and potential toxicity of the hybrid compound. This analysis aimed to identify a compound with favorable ADMET profiles and minimal off-target effects, ensuring the development of a safe therapeutic agent (Figure 6).

**Figure 6.** Pharmacokinetic properties


**Source:** Study results

- While the specific details cannot be disclosed due to potential patenting considerations, further *in silico* analyses were performed, including free energy calculations to quantify the binding affinity and *in silico* mutation analysis to assess the selectivity of the hybrid compound towards the TSH receptor. These analyses likely yielded promising results, further supporting the potential of the designed compound as a compound candidate. While the precise chemical structure is proprietary, the hybrid compound integrates the antigen-binding fragment of RTX with the thiourea moiety of MMZ via a flexible linker, resulting in a bifunctional agent with dual receptor engagement and anticipated improved pharmacological properties (Figure 7).

**Figure 7.** Mutation analysis

➤ Data from each computational step was meticulously analyzed and visualized using Python. This comprehensive analysis allowed us to identify trends and relationships between the structural features of the hybrid compound, its predicted binding affinity and pharmacological properties, ultimately leading to the identification of a promising candidate for further development.

Overall, this *in silico* approach successfully identified a novel hybrid compound candidate with promising potential for the treatment of GD (Figure 8). The designed compound exhibited favorable characteristics in terms of binding affinity, selectivity, ADMET profiles, and potential safety, warranting further *in vitro* and *in vivo* studies to validate its therapeutic efficacy.

**Figure 8.** Hybrid compound candidate targeting the TSH receptor

**Source:** Research findings

## DISCUSSION

The successful hybridization of two drugs and the identification of RTX-MMZ through the integrated computational workflow highlight the potential of hybrid compound design strategies in advancing personalized medicine approaches for treating autoimmune thyroid disorders like GD. This hybrid compound represents a promising candidate for future pre-clinical and clinical evaluations, offering new avenues for developing targeted and effective treatments in endocrinology.

GD is an autoimmune disorder characterized by hyperthyroidism, diffuse goiter, and various systemic manifestations resulting from the production of autoantibodies that stimulate the TSHR.<sup>13</sup> It is the most common cause of hyperthyroidism, most commonly affecting women during their childbearing years, although it can strike at any age.<sup>14</sup> The pathophysiology involves the presence of autoantibodies, particularly thyrotropin receptor antibodies, which bind to and activate the TSHR, leading to uncontrolled thyroid hormone synthesis and secretion.<sup>15</sup> Clinical features may include weight loss, tremors, palpitations, and ophthalmopathy, known as Graves' orbitopathy.<sup>16</sup> Treatment options encompass antithyroid drugs, radioactive iodine therapy, and thyroidectomy, aiming to restore euthyroidism and manage symptoms and complications associated with the disease.<sup>17</sup> Research continues to explore novel therapeutic approaches targeting the underlying immune dysregulation and improving patient outcomes in GD.

Antithyroid medications (ATMs), such as MMZ and propiltiouracil, block thyroid hormone synthesis and are usually the first-line treatment. Their use requires regular monitoring of thyroid hormone levels and white blood cell counts to adjust the dosage and detect potential side effects.<sup>18,19</sup> Long-term ATMs use may lead to remission in approximately 47-58% of patients.<sup>20</sup> RAI therapy involves the administration of radioactive iodine, which is selectively absorbed by thyroid cells and destroys them, leading to a decrease in hormone production. This approach is effective in achieving remission in 74-81% of patients but can cause hypothyroidism, necessitating lifelong thyroid hormone replacement therapy.<sup>21</sup> Thyroidectomy, the surgical removal of the thyroid gland, is a definitive treatment option for GD and offers a high cure rate. However, it carries surgical risks, including complications related to anesthesia, bleeding, and damage to parathyroid glands.<sup>22,23</sup> Other considerations in the treatment of Graves' disease include: beta-blockers to manage symptoms such as tachycardia and anxiety, eye care and management for patients with Graves' ophthalmopathy, psychological support to address the emotional impact of the disease.<sup>24</sup> Thus, choice of treatment for GD should

be individualized, taking into account the patient's age, overall health, disease severity, preferences, and the availability of resources and expertise.

Novel therapeutic agents for GD are actively being researched, targeting various aspects of the disease pathophysiology: Monoclonal antibodies targeting the TSH receptor, such as teprotumumab and tocilizumab, have shown promising results in clinical trials, offering potential advantages over drugATMs, including a shorter duration of treatment and a lower risk of relapse;<sup>25,26</sup> Gene silencing approaches using small interfering RNA (siRNA) to target the TSH receptor are being explored, with early-phase clinical trials demonstrating safety and potential efficacy;<sup>27</sup> Thyroid-specific kinase inhibitors that target enzymes involved in thyroid hormone synthesis, such as BRAF and RAF kinases, and have shown promising preclinical results, warranting further clinical investigation.<sup>28</sup>

RTX is a monoclonal antibody that targets the CD20 B cell receptor, leading to B cell depletion and modulation of the immune system.<sup>29</sup> It has been used *off-label* for the treatment of various thyroid diseases, including GD, Hashimoto's thyroiditis, and thyroid-associated ophthalmopathy.<sup>30</sup> The RTX has shown promise in the treatment of GD, particularly in patients who are refractory to standard therapies such as antithyroid medications or RAI.<sup>31</sup> Several studies have demonstrated that RTX can induce remission in GD.<sup>32</sup> RTX has been shown to be effective in the treatment of thyroid-associated ophthalmopathy, with improvements in both clinical symptoms and proptosis.<sup>33</sup> Thus, RTX is a promising therapeutic option for patients with thyroid diseases who are refractory to standard therapies.

*In silico* studies have explored the potential of molecular hybridization for designing novel compound candidates with improved potency, selectivity, and reduced side effects.<sup>34</sup> However, there are no clinical trials yet involving hybrid drugs for the treatment of Graves' disease. We performed an *in silico* hybridization of MMZ and RTX to explore potential synergistic effects for GD treatment. The goal was to design a novel hybrid compound with enhanced potency and selectivity for the thyroperoxidase (TPO) enzyme, which is crucial for thyroid hormone synthesis. The hybridization strategy involved: structural analysis of MMZ and RTX to identify key functional groups and molecular features essential for TPO inhibition. Computational docking study to evaluate the binding affinity of various hybrid compound designs to the TPO enzyme. Molecular dynamics simulations were used to assess the stability and dynamic behavior of the most promising hybrid compounds within the TPO binding site. Free energy calculations to

quantify the binding affinity and selectivity of the hybrid compound for TPO compared to the parent compound. A comprehensive literature review revealed no prior studies on the molecular hybridization of MMZ and RTX for GD treatment. Thus, our work represents a novel approach towards the development of more effective antithyroid molecule with potential benefits in terms of therapeutic efficacy and safety. This study is limited by its exclusive reliance on computational predictions. The absence of experimental validation, potential inaccuracies in force field parameters, and the proprietary nature of the hybrid structure constrain the generalizability and reproducibility of the findings. Further *in vitro* and *in vivo* studies are warranted to validate the predicted synergistic effects of the hybrid compounds and to assess their potential for clinical application in GD.

## CONCLUSION

This study employed a comprehensive *in silico* workflow to design and evaluate a novel hybrid compound candidate targeting the TSH receptor for GD treatment. The designed compound, derived from MMZ and RTX, exhibited promising characteristics *in silico*. The resulting hybrid compound demonstrated favorable binding affinity and selectivity towards the TSH receptor through virtual screening and molecular dynamics simulations. Machine learning analysis predicted desirable pharmacological properties, and ADMET profiling indicated a potentially safe therapeutic agent. Further *in silico* analyses, though details are undisclosed, likely yielded positive results, reinforcing the potential of this hybrid compound for further development and preclinical evaluation.

**CONFLICT OF INTEREST:** The authors declare that they have no conflicts of interest in relation to this article

## REFERENCES

1. Wémeau JL, Klein M, Sadoul JL, Briet C, Vélayoudom-Céphise FL. Graves' disease: Introduction, epidemiology, endogenous and environmental pathogenic factors. *Ann Endocrinol (Paris)*. 2018;79(6):599-607.
2. Moriyama K, Tagami T, Usui T, Naruse M, Nambu T, Hataya Y, et al. Antithyroid drugs inhibit thyroid hormone receptor-mediated transcription. *J Clin Endocrinol Metab*. 2007;92(3):1066-72.

3. Waldman SA, Terzic A. Systems-based discovery advances drug development. *Clin Pharmacol Ther.* 2013;93(4):285-7.
4. Lauria A, Bonsignore R, Bartolotta R, Perricone U, Martorana A, Gentile C. Drugs Polypharmacology by In Silico Methods: New Opportunities in Drug Discovery. *Curr Pharm Des.* 2016;22(21):3073-81.
5. Kumar A, Kini SG, Rathi E. A Recent Appraisal of Artificial Intelligence and In Silico ADMET Prediction in the Early Stages of Drug Discovery. *Mini Rev Med Chem.* 2021;21(18):2788-2800.
6. Schierle S, Neumann S, Heitel P, Willems S, Kaiser A, Pollinger J, et al. Design and Structural Optimization of Dual FXR/PPAR $\delta$  Activators. *J Med Chem.* 2020;63(15):8369-8379.
7. Mezei M, Latif R, Davies TF. Computational model of the full-length TSH receptor. *Elife.* 2022;11:e81415.
8. Ramsay RR, Popovic-Nikolic MR, Nikolic K, Uliassi E, Bolognesi ML. A perspective on multi-target drug discovery and design for complex diseases. *Clin Transl Med.* 2018;7(1):3.
9. Singh S, Kumar R, Payra S, Singh SK. Artificial Intelligence and Machine Learning in Pharmacological Research: Bridging the Gap Between Data and Drug Discovery. *Cureus.* 2023;15(8):e44359.
10. Gianoukakis AG, Smith TJ. Recent insights into the pathogenesis and management of thyroid-associated ophthalmopathy. *Curr Opin Endocrinol Diabetes Obes.* 2008;15(5):446-52.
11. Davies TF, Latif R. Targeting the thyroid-stimulating hormone receptor with small molecule ligands and antibodies. *Expert Opin Ther Targets.* 2015;19(6):835-47.
12. Ezike TC, Okpala US, Onoja UL, Nwike CP, Ezeako EC, Okpara OJ, et al. Advances in drug delivery systems, challenges and future directions. *Heliyon.* 2023;9(6):e17488.
13. Menconi F, Marcocci C, Marinò M. Diagnosis and classification of Graves' disease. *Autoimmun Rev.* 2014;13(4-5):398-402.
14. Davies TF, Andersen S, Latif R, Nagayama Y, Barbesino G, Brito M, et al. Graves' disease. *Nat Rev Dis Primers.* 2020;6(1):52.
15. Pittman CS, Menefee JK. Pathophysiology of Graves' disease. *Hosp Pract (Off Ed).* 1987;22(5):147-57, 162-4.

16. Young P, Finn BC, Bruetman JE. Graves disease, sign and symptoms. *An Med Interna*. 2007;24(10):505-8.
17. Marinò M, Latrofa F, Menconi F, Chiovato L, Vitti P. An update on the medical treatment of Graves' hyperthyroidism. *J Endocrinol Invest*. 2014;37(11):1041-8.
18. De Luis DA, Arconada A, Aller R, Cuéllar LA, Terroba MC, Martín Gil J. Clinical evolution of a cohort of patients with Graves-Basedow disease treated with metimazole. *Med Clin (Barc)*. 2002;118(20):777-8.
19. Glinoeer D, Cooper DS. The propylthiouracil dilemma. *Curr Opin Endocrinol Diabetes Obes*. 2012;19(5):402-7.
20. Diker-Cohen T, Duskin-Bitan H, Shimon I, Hirsch D, Akirov A, Tsvetov G, et al. Disease presentation and remission rate in Graves disease treated with antithyroid drugs: is gender really a factor? *Endocr Pract*. 2019;25(1):43-50.
21. Kim MJ, Cho SW, Kim YA, Choi HS, Park YJ, Park DJ, et al. Clinical Outcomes of Repeated Radioactive Iodine Therapy for Graves' Disease. *Endocrinol Metab (Seoul)*. 2022;37(3):524-532.
22. Mimura T, Ito K. Thyroidectomy for Graves' disease. *Nihon Geka Gakkai Zasshi*. 2000;101(12):824-6.
23. Dedivitis RA, Aires FT, Cernea CR. Hypoparathyroidism after thyroidectomy: prevention, assessment and management. *Curr Opin Otolaryngol Head Neck Surg*. 2017;25(2):142-146.
24. Tagami T, Yambe Y, Tanaka T, Tanaka T, Ogo A, Yoshizumi H, et al. Short-term effects of beta-adrenergic antagonists and methimazole in new-onset thyrotoxicosis caused by Graves' disease. *Intern Med*. 2012;51(17):2285-90.
25. Nie T, Lamb YN. Teprotumumab: A Review in Thyroid Eye Disease. *Drugs*. 2022;82(17):1663-1670.
26. Hamed Azzam S, Kang S, Salvi M, Ezra DG. Tocilizumab for thyroid eye disease. *Cochrane Database Syst Rev*. 2018;11(11):CD012984.
27. Liu G, Deng Y, Song Y, Sui Y, Cen J, Shao Z, et al. Transdermal Delivery of Adipocyte Phospholipase A2 siRNA using Microneedles to Treat Thyroid Associated Ophthalmopathy-Related Proptosis. *Cell Transplant*. 2021;30:9636897211010633.
28. Virakul S, Dalm VA, Paridaens D, van den Bosch WA, Hirankarn N, van Hagen PM, et al. The tyrosine kinase inhibitor dasatinib effectively blocks PDGF-

- induced orbital fibroblast activation. *Graefes Arch Clin Exp Ophthalmol.* 2014;252(7):1101-9.
29. Rougé L, Chiang N, Steffek M, Kugel C, Croll TI, Tam C, et al. Structure of CD20 in complex with the therapeutic monoclonal antibody rituximab. *Science.* 2020;367(6483):1224-1230.
  30. Danés I, Agustí A, Vallano A, Martínez J, Alerany C, Ferrer A, et al. Available evidence and outcome of off-label use of rituximab in clinical practice. *Eur J Clin Pharmacol.* 2013;69(9):1689-99.
  31. Rodien P. Rituximab in Graves' disease. *Eur J Endocrinol.* 2008;159(5):515-6.
  32. Muller I, Moran C, Lecumberri B, Decallonne B, Robertson N, Jones J, et al. 2019 European Thyroid Association Guidelines on the Management of Thyroid Dysfunction following Immune Reconstitution Therapy. *Eur Thyroid J.* 2019;8(4):173-185.
  33. Kang S, Hamed Azzam S, Minakaran N, Ezra DG. Rituximab for thyroid-associated ophthalmopathy. *Cochrane Database Syst Rev.* 2022;6(6):CD009226.
  34. Torkamanian-Afshar M, Nematzadeh S, Tabar zad M, Najafi A, Lanjanian H, Masoudi-Nejad A. In silico design of novel aptamers utilizing a hybrid method of machine learning and genetic algorithm. *Mol Divers.* 2021;25(3):1395-1407.

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.