SciELO Preprints

Publication status: This preprint has not been published elsewhere.

Trends and extreme events in air temperature and precipitation in Goiás, Brazil

Guilherme Cardoso da Silva, Diego Tarley Ferreira Nascimento

https://doi.org/10.1590/SciELOPreprints.16770

Submitted on: 2026-07-02
Posted on: 2026-11-03 (version 2)
(YYYY-MM-DD)

Version justification: Correção da Referência Bibliográfica.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

TRENDS AND EXTREME EVENTS IN AIR TEMPERATURE AND PRECIPITATION IN GOIÁS, BRAZIL

Guilherme Cardoso da Silva

Federal University of Goiás, Goiânia, Goiás, Brazil.

ORCID: https://orcid.org/0009-0006-7445-1447

Diego Tarley Ferreira Nascimento

Professor Federal University of Goiás, Goiânia, Goiás, Brazil.

Brazil.ORCID: https://orcid.org/0000-0002-0420-3636

ABSTRACT

Climate change is unequivocal, widespread, accelerating, irreversible and without historical precedent, with extreme climatic events recorded in all regions of the planet, causing adverse impacts in the environmental, economic and social spheres. The aim of this research was to detect climate change trends and identify extreme events and in the The study aimed to identify climate trends and extreme weather events in the state of Goiás, using air temperature and precipitation indices from a 45-year time series (1980 to 2024), based on daily data recorded by seven conventional weather stations of the National Meteorological Institute, located in Goiânia, Jataí, Pirenópolis, Aragarças, Formosa, Posse and Ipameri. To this end, indices developed by the Expert Team on Climate Change Detection and Indices (ETCCDI), linked to the World Meteorological Organization (WMO), were used. Seventeen indices were calculated with the R software, using the Rclimdex extension. For the precipitation-related indices, the data show a negative trend with high statistical significance in only a few stations. In the case of the temperature-related indices, there is a positive trend with high statistical significance at all the stations analyzed, which shows an increase in maximum and minimum temperatures over the time series analyzed. This study demonstrates significant warming trends and increasing climate extremes in Goiás, reinforcing the need for evidence-based adaptation and mitigation measures to enhance resilience and promote sustainable development under a changing climate.

Keywords: Climate Change, Global Warming, Climate Trends, Climate Indices, Rclimdex.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

INTRODUCTION

Understanding the variability and changes in climatic parameters is of paramount importance, as it forms the basis for environmental diagnostics and prognostics. Such insights are critical for guiding the planning of human activities and mitigating or preventing undesirable losses and damages.

The Intergovernmental Panel on Climate Change (IPCC, 2021) is categorical in stating that global climate change is a reflection of human actions, pointing out that these changes are widespread, accelerated, irreversible and without historical precedent. The same organism further underscores the increasing frequency, intensity, and severity of extreme weather events worldwide, often associated with extreme temperature anomalies (e.g., heat or cold waves), intense rainfall episodes, or prolonged droughts.

Recent decades have witnessed record-breaking heatwaves in Europe, Asia, North America, and other regions, as well as increasingly severe precipitation events, prolonged droughts, and compound extremes affecting ecosystems, infrastructure, food production, water resources, and human health.

Among the most significant recent examples are the extreme heatwaves recorded in Europe, China, and the United States during 2023, when temperatures exceeded historical records in several regions – including the registration of 52.2°C in Sanbao (Xinjiang, China) – according to the China Meteorological Administration (CMA). Initially, 2023 was declared the hottest year on record since measurements began in 1850 (WMO, 2024). However, 2024 surpassed this record, becoming the first year to exceed 1.5°C above pre-industrial levels (COPERNICUS, 2025).

At the same time, extreme precipitation events have become more frequent and destructive in many parts of the world, causing floods, landslides, and substantial economic losses. Conversely, prolonged droughts have affected extensive areas of South America, Africa, Europe, and western North America, compromising water availability, agricultural production, and energy generation.

In South America, and particularly in Brazil, the effects of climate variability and change have become increasingly evident through both temperature and precipitation extremes. In the winter of 2022, a cold air mass led to record-low

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

temperatures in the Midwest and Southeast regions of Brazil, with minimum temperatures recorded at 4.9°C in Brasília (DF), 7.0°C in Cuiabá (MT), 7.6°C in Belo Horizonte (MG), and 7.8°C in Goiânia (GO). In turn, a heatwave in 2023 resulted in record-high temperatures, including 44.8°C in Araçuaí (MG) and 44.2°C in Cuiabá (MT), according to data from the National Institute of Meteorology (INMET).

Also, notable examples include the extreme rainfall events in Northeast and Southeast Brazil in 2022 and in Rio Grande do Sul in 2024, which resulted in hundreds of fatalities and displaced thousands of individuals. In contrast, severe droughts have affected extensive areas of the North, Northeast, and Central-West regions, notably during the Amazon droughts of 2005, 2010, and 2023–2024, the prolonged Northeast drought between 2012 and 2017, and the widespread drought that impacted Central Brazil during 2020–2021, causing substantial impacts on water resources, agriculture, energy production, navigation, and ecosystem functioning.

While extreme weather events are not novel phenomena unique to recent years in Brazil (ARTAXO, 2014, 2020; MARENGO, 2009; MARENGO, CARDONA, MARTINEZ, 2022), the current global climate crisis is characterized by their increased recurrence, intensity, duration, and severity. These events disproportionately affect countries, communities, and ecosystems, with greater impacts on those that are more vulnerable or sensitive (MARENGO et al., 2009).

In this context, the Central-West Region of Brazil warrants particular attention due to its economic reliance on agriculture and livestock. These sectors are highly susceptible to fluctuations in temperature and precipitation extremes, which can lead to reduced agricultural productivity, crop failures, and livestock mortality (CASSOL; BOHNER, 2012). Beyond economic losses, climatic irregularities and anomalies also impact water availability, affecting food production, domestic water supply, and hydroelectric power generation. This, in turn, exacerbates challenges related to food, water, and energy security (RODRIGUES, 2017).

The IPCC (2021) projects that the Brazilian Midwest will experience a trend toward higher temperatures, reduced relative humidity, and altered precipitation patterns. This projection is supported by scientific studies conducted in the state of

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Goiás (NEVES, 2018; DEUS; NASCIMENTO, 2021) and the broader Central-West Region (ALMEIDA, 2012; SANTOS et al., 2023). Thus, the general objective of this study is to identify climate trends and extreme weather events associated with air temperature and precipitation indices across seven weather stations in the state of Goiás, using a 45-year time series (1980–2024). The analysis employs standardized indices developed by the Expert Team on Climate Change Detection and Indices (ETCCDI), affiliated with the World Meteorological Organization (WMO).

It is hoped that this work will corroborate the climate change scenarios projected by the IPCC for the state of Goiás and elucidate trends and occurrences of climatic extremes in the seven selected locations, which represent diverse climatic typologies: Goiânia, Jataí, Pirenópolis, Aragarças, Formosa, Posse, and Ipameri. The findings aim to provide a scientific foundation for the formulation of public policies and strategic planning to address the climate crisis in the region.

METHODOLOGY

LOCATION AND BRIEF CHARACTERIZATION OF THE STUDY AREA

The area under study refers to the state of Goiás, located in the Central-West Region of Brazil, covering an area of 340,242 km² - which represents approximately 4% of the national territory. Specifically, the study focuses on seven municipalities within Goiás where conventional weather stations are situated. These stations were selected due to their extensive and consistent time series data: Goiânia, Jataí, Pirenópolis, Aragarças, Formosa, Posse, and Ipameri (Figure 1).

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770




Figure 1 – Location, climate types, and selected stations in the state of Goiás. Source: authors, according to the Koppen-Geiger (1961) climate classification system.

As noted by Nimer (1979), the study region exhibits an average annual temperature of 22°C and average of annual precipitation between 1,550 mm and 1,750 mm, with a dry season lasting four months, typically from May to August. In turn, based on 105 ANA rainfall stations and 47 INMET climate stations, Cardoso, Marcuzzo, and Barros (2014) report an average annual temperature of 23.4°C for the area in question, with higher values in the northwest region, reaching an average of 27°C. The authors report an average annual precipitation of 1,500 mm, ranging from 1,000 to 2,800 mm.

According to the most recent data, based on the 1991-2020 climate norms calculated by INMET for the stations used in the study (Table 1), the highest average annual temperature is recorded in Aragarças (26°C), and the lowest in Formosa and Jataí (22.8°C) – due to the influence of the continental climate, altitude, and latitude, respectively.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Station	Average Annual Temperature (°C)	Lowest monthly average minimum temperature (°C)	Highest monthly average maximum temperature (°C)	Average Annual Precipitation (mm)	Length of the dry season (months)
Posse	24.4	17.8 (julho)	32.5 (setembro)	1,345.5	5
Formosa	22.8	14.6 (julho)	30.8 (setembro)	1,339.4	5
Aragarças	26	15.5 (julho)	36.5 (setembro)	1,450.9	5
Pirenópolis	23.3	14.2 (julho)	33.7 (setembro)	1,725.9	5
Goiânia	24.5	14.8 (julho)	34.0 (setembro)	1,667.3	5
Jataí	22.8	11.6 (julho)	33.4 (setembro)	1,593.3	3
Ipameri	22.9	13.1 (julho)	31.8 (outubro)	1,481.8	5

Table 1 – Annual and monthly averages of mean, maximum, and minimum temperatures, as well as precipitation and the duration of the dry season. Source: climate norms calculated by INMET (1991/2020).

As illustrated in Figure 1, the predominant climate typology in Goiás is a hot and semi-humid tropical climate, characterized by average temperatures exceeding 18°C throughout the year and a dry period of 4 to 5 months – according to the Koppen-Geiger (1961) climate classification system. This condition is reflected in the weather stations located in Posse, Formosa, Aragarças, Pirenópolis, and Goiânia. Additionally, a hot and humid tropical climate typology is observed in the study area, with only 3 dry months, as exemplified by the Jataí station.

Furthermore, a sub-hot tropical climate is present in the region, characterized by average temperatures between 15°C and 18°C in at least one month of the year. In the study area, this typology is primarily associated with semi-humid conditions (with 5 dry months), as represented by the Ipameri meteorological station. However, the sub-hot tropical climate can also occur under humid conditions (only 3 dry months), although this subtype is not represented by any of the selected stations in this investigation.

According to Campos et al. (2002), Luiz (2012), Neves (2018; 2019), and Nascimento and Oliveira (2020), the region's climate is influenced by physiographic characteristics and the interaction of atmospheric systems of equatorial and tropical origin, with occasional incursions of polar systems. These dynamics result in two well-defined seasons: a hot and rainy spring/summer, driven by the

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

dominance of the continental Equatorial mass and the South Atlantic Convergence Zone, and a milder and drier autumn/winter, influenced by the Atlantic Tropical mass and continentalized polar air incursions.

METHODOLOGICAL PROCEDURES

To conduct this study, daily precipitation and air temperature data (maximum and minimum) were obtained from seven conventional weather stations managed by the National Institute of Meteorology (INMET) and located in the state of Goiás (Formosa, Posse, Aragarças, Pirenópolis, Goiânia, Jataí, and Ipameri).

A common time series spanning 45 years (1980–2024) was used for all stations, corresponding to 16,437 daily records. Although conventional weather stations typically provide longer and more consistent data series, some gaps were identified. The maximum proportion of missing data was 10.6% for maximum temperature (Ipameri), 7.6% for minimum temperature (Formosa), and 2.8% for precipitation (Aragarças). These percentages are within the range generally considered acceptable in climatological studies (WMO, 2017).

In addition, the calculation of climate indices allows missing observations to be coded as −99.9 and includes a quality-control procedure to assess the potential influence of missing data on the results (ZHANG; YANG, 2004). So, considering the low proportion of missing values, no data infilling procedures were applied in order to preserve the original variability of the series and avoid artificial homogenization that could affect the derived climate indices.

To detect climate trends and extreme weather events, indices developed by the Expert Team on Climate Change Detection and Indices (ETCCDI), affiliated with the World Meteorological Organization (WMO), were utilized. These indices are designed to measure and detect variability and changes in climate parameters through standardized metrics, enabling global-scale observation and comparison (KARL, NICHOLLS, GHAZI, 1999). A comprehensive description of the indices can be found in the full workshop report (PETERSON et al., 2001) or its summary (PETERSON, 2005).

The standardized ETCCDI/WMO indices are widely used in scientific literature at various scales. In Brazil, studies have been conducted for the entire country

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

(SALVIANO; GROPPO; PELLEGRINO, 2016; PENEREIRO; MESCHIATTI, 2018; REGOTO, et al., 2021; COELHO et al., 2023), as well as for specific biomes LMEIDA, et al., 2016; JOSEPH. SOUZA. SABINO, 2021; SILVA et al., 2019; LUIZ-SILVA et al., 2025), regions (ALMEIDA, 2012; COSTA et al., 2020; SANTOS et al., 2023; SANTOS et al., 2024), and Federal Units (REBELLO, et al., 2008; SANTOS, et al., 2009; SILVA et al., 2015; SILVA, MONTENEGRO, SOUZA, 2017; CASTELHANO; PINTO, 2022; LUIZ-SILVA; OSCAR-JUNIOR, 2022), in addition to numerous local-scale studies.

Among the 27 indices proposed by ETCCDI/WMO, 19 were selected for this study: 11 related to air temperature and 8 to precipitation, as detailed in Table 2. The data were organized in Excel spreadsheets and processed using the Rclimdex extension in the R program, following the recommendations of Zhang and Yang (2004) and Zhang et al. (2005).

To assess statistical significance, linear trends are calculated using the least squares method (slope) were considered significant if they exceeded the standard error of estimate (STD). Trends with p-values below 0.05, as determined by the Student's t-test, were classified as highly statistically significant, while those with p-values between 0.05 and 0.1 were considered moderately significant, following the approach suggested by Santos et al. (2009).

It is worth noting that, although linear regression based on least squares is the standard approach adopted in the ETCCDI/RClimDex framework, it can be sensitive to outliers, deviations from normality, and serial correlation. Therefore, future studies could complement the analysis by using nonparametric approaches, such as the Mann-Kendall test and the Theil-Sen slope estimator, due to their greater robustness to non-normality, autocorrelation, and outliers.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Acronym	Name	Description	Unit
TXx	Highest Maximum Temperature	Maximum value of daily maximum temperature	°C
TXn	Lowest Maximum Temperature	Minimum value of daily maximum temperature	°C
TNx	Highest Minimum Temperature	Maximum value of daily minimum temperature	°C
TNn	Lowest minimum temperature	Minimum value of daily minimum temperature	°C
TX10p	Cold days	Percentage of days in which the maximum daily temperature was below the 10th percentile	%
TX90p	Hot days	Percentage of days in which the maximum daily temperature was above the 90th percentile	%
TN10p	Cold nights	Percentage of days in which the minimum daily temperature was below the 10th percentile	%
TN90p	Hot nights	Percentage of days in which the minimum daily temperature was above the 90th percentile	%
WSDI	Duration of hot periods	Annual count of days with at least 6 consecutive days in which the maximum daily temperature was above the 90th percentile	days
CSDI	Duration of cold periods	Annual count of days with at least 6 consecutive days in which the minimum daily temperature was below the 10th percentile	days
DTR	Diurnal temperature range	difference between temperature maximum and minimum	°C
RX1day	Maximum amount of precipitation in 1 day	Maximum volume of precipitation in one day	mm
RX5day	Maximum amount of precipitation in 5 days	Maximum precipitation volume in five consecutive days	mm
R10mm	Days with heavy precipitation (>10mm)	Number of days in a year with precipitation greater than 10 mm	days
R20mm	Days with intense precipitation (>20mm)	Number of days in a year with precipitation greater than 20 mm	days
R40mm	Days with very intense precipitation (>40mm)	Number of days in a year with precipitation greater than 40 mm	days
CDD	Consecutive Dry Days	Maximum number of consecutive days with precipitation less than 1 mm	days
CWD	Consecutive Wet Days	Maximum number of consecutive days with precipitation greater than 1 mm	days
PRCPTOT	Total Precipitation	Total annual precipitation	mm

Table 2 – List of climate indices used. Source: authors.

RESULTS AND DISCUSSIONS

The values of the climate trend indices for the seven selected stations in the state of Goiás, based on the period from 1980 to 2024, are presented in Table 3. For the temperature-related indices, a predominance of positive trends (indicating an increase) was observed, with high statistical significance across all stations for the highest and lowest maximum temperatures (TXx and TXn), highest and lowest minimum temperatures (TNx and TNn), and diurnal temperature range (DTR).

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Indices / Stations		Aragarças	Pirenópolis	Goiânia	Posse	Formosa	Jataí	Ipameri
TXx	Highest Max. Temperature	**0.097**	**0.102**	**0.107**	**0.088**	**0.069**	**0.079**	**0.098**
TXn	Lowest Max. Temperature	**0.048**	**0.084**	**0.074**	**0.024**	**0.066**	**0.052**	**0.09**
TNx	Highest Min. Temperature	**0.041**	**0.040**	**0.074**	**0.056**	**0.08**	**0.037**	**0.045**
TNn	Lowest Min. temperature	**0.015**	**0.018**	**0.058**	**0.009**	**0.034**	**0.005**	**-0.007**
TX10p	Cold days	-0.369	-0.633	-0.416	-0.426	-0.412	**-0.232**	-0.32
TX90p	Hot days	0.692	0.889	0.802	0.607	0.57	0.599	0.646
TN10p	Cold nights	0.087	-0.299	-0.564	-0.187	-0.349	-0.314	-0.298
TN90p	Hot nights	0.130	0.495	0.625	**0.352**	0.701	0.471	0.462
WSDI	Duration of hot periods	0.424	0.598	0.811	0.309	0.36	0.630	0.559
CSDI	Duration of cold periods	-0.066	**-0.072**	-0.483	-0.227	-0.223	-0.068	-0.084
DTR	Diurnal temperature range	**0.055**	**0.047**	**0.010**	**0.033**	**0.013**	**0.016**	**0.036**
RX1day	Precipitation in 1 day	0.103	-0.256	0.283	-0.421	0.032	0.054	0.171
RX5day	Precipitation in 5 days	-0.559	-0.829	0.056	-0.512	-0.871	-0.292	0.168
R10mm	Days with heavy precip.	-0.174	-0.187	-0.122	-0.299	-0.279	-0.197	-0.191
R20mm	Days with intense precip.	-0.071	-0.021	-0.058	-0.188	-0.141	-0.093	-0.006
R40mm	Days with very intense precip	**-0.030**	**0.000**	**0.031**	**-0.067**	**-0.036**	**0.016**	**-0.022**
CDD	Consecutive Dry Days	0.630	0.574	1.132	0.543	0.795	0.321	0.238
CWD	Consecutive Wet Days	**0.002**	**0.008**	**-0.001**	**-0.060**	**-0.075**	**-0.058**	**-0.093**
PRCPTOT	Total Precipitation	**-6.620**	-2.849	**-0.720**	**-8.786**	**-7.656**	**-4.703**	**-3.551**

Table 3 – Trend detection indices for selected stations in Goiás for the period from 1980 to 2024. Source: authors, based on data stations and calculations carried out in RClimDEX. Note: positive trends in green and negative trends in red, and high statistical significance in dark gray (p-value < 0.05) and good statistical significance in light gray (p-value between 0.05 and 0.1).

The indices for hot days (TX90p) and hot nights (TN90p) also exhibited positive trends (increase), with good statistical significance in most locations, except for the Aragarças and Formosa stations, where no significance was observed. The hot period duration index (WSDI) showed a positive trend but lacked statistical significance in all studied locations. These findings align with the global scenario of rising temperatures (IPCC, 2021) and are consistent with previous studies for the state of Goiás (Almeida, 2012; Neves, 2018; Hofmann et al., 2021, 2023; Santos et al., 2023; Silva and Nascimento, 2021), Brazil (Regoto et al., 2021), and the South American continent (Skansi et al., 2013).

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Conversely, the indices for cold days (TX10p), cold nights (TN10p), and cold periodo duration (CSDI) displayed negative trends (decrease), with good statistical significance in most locations. This suggests a reduction in the frequency of colder days and nights, consistent with findings by Santos et al. (2006), Nascimento, Santos and Barros (2011), and Luiz (2012) for Goiânia, and Almeida (2012) and Santos et al. (2023) for the broader Central-West Region.

Regarding precipitation-related indices, negative trends (reduction) predominated for the maximum 5-day precipitation (RX5day), days with precipitation exceeding 10 mm (R10mm), 20 mm (R20mm), and 40 mm (R40mm), consecutive wet days (CWD), and total annual precipitation (PRCPTOT). The R40mm, CWD, and PRCPTOT indices showed high statistical significance, while R10mm and R20mm exhibited good significance across all stations. In contrast, the consecutive dry days (CDD) index showed a positive trend at all stations, and the maximum 1-day precipitation (RX1day) index increased in five of the seven locations, though neither trend was statistically significant.

These precipitation trends are consistent with findings by Alexander et al. (2006) for much of the globe, Salviano, Groppo, and Pellegrino (2016) for over 70% of Brazil's territory, and Santos et al. (2006), Penereiro and Meschiatti (2018), Santos et al. (2023), and Neves et al. (2024) for the state of Goiás. The results indicate a concerning scenario: a reduction in annual precipitation totals, fewer days with rainfall (whether consecutive or not), and a shorter wet season. This aligns with IPCC projections for the Central-West Region and is supported by studies such as Skansi et al. (2013), Almeida (2012), Marcuzzo, Oliveira and Cardoso (2013), Neves (2018), and Santos et al. (2023).

Analysis of the time series based on ETCCDI/WMO indices also highlights extreme temperature events. As shown in Figure 2, the highest maximum temperature (TXx) recorded was 43.3°C at the Aragarças station in October 2020, followed by 42.0°C in Jataí and 41.2°C in Goiânia during the same month and year. These values exceed previous records for the study area (NASCIMENTO, LIMA, CRUZ, 2019; DEUS, NASCIMENTO, 2021), underscoring the breaking of temperature records year after year. Extreme maximum temperatures were also recorded in 2024 at the Pirenópolis and Posse stations, a year marked as the

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

hottest since records began and the first to exceed 1.5°C above pre-industrial levels (COPERNICUS, 2025).




Figure 2 – Spatial variation of extreme events from 1980 to 2024 for some indices associated with air temperature (TXx, TNn, WSDI and DTR). Source: authors, based on INMET data processed in RClimDEX.

The lowest minimum temperature (TNn) records were -1.0°C in 1988 at Jataí, 0.6°C in 1981 at Ipameri, and 2.6°C in 1994 in Goiânia. The first and last locations, situated in the southwest and southeast portions of Goiás, respectively, are more susceptible to incursions of polar air masses, as noted by Nascimento (2016) and Neves (2019). Such low temperatures pose significant risks to agricultural crops, livestock, and human health, particularly for vulnerable populations such as the elderly, children, and those with chronic illnesses (FENNER, SILVA, GURGEL, 2009; MENDONÇA, 2005).

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Heatwaves were another notable extreme event, with a record duration of 96 days in Goiânia in 2024. Similar records were observed in Jataí (89 days), Aragarças (53 days), Pirenópolis (57 days), and Ipameri (46 days) during the same year. These prolonged heatwaves, attributed to the influence of the El Niño phenomenon, pose serious health risks, including increased cardiovascular and respiratory mortality, particularly among vulnerable groups such as the elderly, pregnant women, and individuals with chronic conditions (MASSELOT et al., 2023; ROMANELLO et al., 2024). Furthermore, there is a great risk of dehydration, especially as the period is dry and associated with lower levels of relative humidity.

High diurnal temperature ranges (DTR) were also observed, particularly at the end of winter and beginning of spring. Record DTR values included 21.7°C in Goiânia and Jataí and 21.6°C in Aragarças, recorded in August of 1994, 2023, and 2014, respectively. Such fluctuations can exacerbate health risks and impact agricultural productivity.

The maps in Figure 3 illustrate extreme values of maximum 1-day precipitation (RX1day), with notable records of 174 mm in Posse (1994), 167 mm in Pirenópolis (1989), and 151 mm in Jataí (2005). Even more alarming were the extreme 5-day precipitation events, with records of 390 mm in Pirenópolis, 358 mm in Aragarças, and 245 mm in Goiânia.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770




Figure 3 – Spatial variation of extreme events from from 1980 to 2024 for some indices associated with precipitation (RX1day, RX5day, CDD and PRCPTOT). Source: authors, based on INMET data processed in RClimDEX.

These intense precipitation events have led to recurring urban flooding, as highlighted by Rego and Barros (2014) and Luiz and Romão (2019) for Goiânia (Figure 4-A). Deus and Nascimento (2016) also documented the impacts of extreme rainfall in smaller cities, such as Pirenópolis (Figure 4-B), a tourist destination in Goiás.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770



Figure 4 – Flooding recorded in the cities of Goiânia (A) and Pirenópolis (B) following extreme rainfall events. Source: (A): https://opopular.com.br/cidades/chuva-forte-causa-estragos-em-goiania-veja-fotos-e-video-1.3125219 and (B): https://opopular.com.br/cidades/defesa-civil-de-pirenopolis-alerta-para-risco-de-desabamentos-e-recomenda-fechamento-de-cachoeiras-1.2384903

The occurrence of extreme precipitation events in the study area is often linked to the performance and intensity of atmospheric systems, such as the South Atlantic Convergence Zone and the continental equatorial mass, influenced by atmospheric teleconnections like the El Niño Southern Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO) (Nascimento, Lima, & Cruz, 2019; Silva & Carpenedo, 2021; Alves, Cabral, & Nascimento, 2022).

The maximum annual value of consecutive dry days (CDD) reached 177 days in Posse and Formosa, followed by 159 days in Pirenópolis and 156 days in Goiânia. Such prolonged dry periods, lasting up to five or six months, severely impact water, food, and energy security (Hoff, 2011; Mariani et al., 2016). Annual precipitation (PRCPTOT) records included 2,437 mm in Aragarças (1983), 2,417 mm in Pirenópolis (1989), and 2,193 mm in Goiânia (2021).

The northeastern portion of Goiás, where Posse and Formosa are located, historically receives lower precipitation due to topographic barriers that hinder the

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

entry of rain-producing atmospheric systems (NASCIMENTO, 2016; NEVES, 2019). In contrast, Aragarças, Pirenópolis, and Goiânia are situated in areas less affected by such barriers.

The dry season in the study area is primarily influenced by the dominance of the Atlantic Tropical Mass during autumn and winter, which brings high temperatures, atmospheric stability, and a lack of precipitation (SERRA; RATISBONA, 1942; MONTEIRO, 1951). However, in particularly dry years, the absence of rainfall for three or more consecutive months is common, often exacerbated by atmospheric teleconnections such as ENSO and PDO (NASCIMENTO; COSTA, 2020; ALVES, CABRAL, NASCIMENTO, 2022).

The results of this study align with the broader scientific literature and IPCC projections, indicating a hotter and drier climate for the study area, with increased intensity and recurrence of extreme temperature and precipitation events. These findings underscore the urgent need for adaptive strategies to mitigate the impacts of climate change on agriculture, water resources, and public health in the state of Goiás.

CONCLUSIONS

The application of climate indices has proven highly effective in identifying trends in meteorological variables across selected weather stations in the state of Goiás. The analysis revealed a clear and statistically significant pattern of increasing temperatures, as evidenced by positive trends in the highest and lowest maximum and minimum temperatures (TXx, TXn, TNx, and TNn) and the diurnal temperature range (DTR). These findings align with global observations of rising temperatures and underscore the intensification of warming trends in the region.

In contrast, precipitation-related indices predominantly exhibited negative trends, particularly in the number of days with heavy (R10mm), intense (R20mm), and very intense (R40mm) rainfall, consecutive wet days (CWD), and total annual precipitation (PRCPTOT). While the trends for PRCPTOT, CWD, and R40mm were statistically significant, other indices showed good but not high significance. This suggests a shift toward drier conditions, albeit with variability across the study area.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

The study also highlighted extreme climatic events recorded over the 45-year period (1980–2024), including: a highest maximum temperature of 44.3°C in Aragarças (2020); a lowest minimum temperature of -1.0°C in Jataí (1988); a maximum 1-day precipitation of 174 mm in Posse (1999), a record of 177 consecutive dry days in Posse and Formosa, and a total annual precipitation of 2,437.9 mm in Aragarças (1983).

These findings are consistent with the Sixth Assessment Report (AR6) of the Intergovernmental Panel on Climate Change (IPCC, 2021), which emphasizes unprecedented global warming and the increased frequency and intensity of extreme weather events. For the Central-West Region of Brazil, the IPCC projects further temperature increases and potential alterations in precipitation patterns, though the latter lacks consistent statistical significance in this study.

While this study provides valuable insights into climate trends and extremes in Goiás, further research is needed to deepen our understanding of these phenomena. Future studies should a) conduct more granular analyses of specific indices to identify localized trends and impacts; b) investigate the socioeconomic and environmental repercussions of rising temperatures and changing precipitation patterns; and c) explore the role of atmospheric teleconnections, such as ENSO and PDO, in driving extreme events in the region.

Finally, the findings of this study underscore the urgent need for proactive measures to address the impacts of climate change in Goiás. By integrating scientific knowledge with practical adaptation and mitigation strategies, the state can enhance its resilience to climate variability and ensure sustainable development for future generations.

ACKNOWLEDGEMENTS

The authors would like to thank the Coordination for the Improvement of Higher Education Personnel (CAPES), for awarding a master's degree scholarship to the first author, and the National Council for Scientific and Technological Development (CNPq), for awarding a research productivity scholarship and for the financial support under Public Call MCTI/CNPq No. 14/2023 to the second author. The authors would also like to thank the Pro-Rectory of Extension and Culture of the

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Federal University of Goiás for the financial support for translation and proofreading services.

AUTHOR'S CONTRIBUTIONS

Silva, G.C. - The author contributed to the elaboration, realization and manipulation of the data and writing.

Nascimento, D.T.F. - The author contributed to the elaboration, realization and manipulation of the data and writing.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

RESEARCH DATA AVAILABILITY

All the data supporting the results of this study were published in the article itself.

USE OF AI

No Artificial Intelligence tools were used.

BIBLIOGRAPHIC REFERENCES

ALEXANDER, L. V.; et al. Global Observed Changes in Daily Climate Extremes of Temperature and Precipitation. Journal of Geophysical Research, v. 111, D05109, 2006.

ALMEIDA, J. M. V. de. Índices de monitoramento e detecção de mudanças climáticas na região Centro-Oeste do Brasil. 2012. 178f. Dissertação (Mestrado em Meteorologia) – Pós-Graduação em Meteorologia, Centro de Tecnologia e Recursos Naturais, Universidade Federal de Campina Grande, Paraíba, Brasil, 2012.

ALVES, W. S.; CABRAL, J. B. P.; NASCIMENTO, D. T. F. A participação do El Niño Oscilação Sul (ENOS) e da Oscilação Decadal do Pacífico (ODP) nas anomalias de chuva em Goiás e no Distrito Federal. Revista Ra'eGa - Espaço Geográfico em Análise, v. 55, p. 176-200, 2022. http://dx.doi.org/10.5380/raega.v55i0.83567.

ARTAXO, P. As três emergências que nossa sociedade enfrenta: saúde, biodiversidade e mudanças climáticas. Estudos Avançados, v. 34, p. 53-66, 2020.

ARTAXO, P. Mudanças climáticas e o Brasil. Revista USP, [S. l.], n. 103, p. 8-12, 2014.

CAMPOS, A. B. de. et al. Análise do comportamento espacial e temporal das temperaturas e pluviosidades no Estado de Goiás. In: ALMEIDA, M. G. de (Org.).

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

Abordagens geográficas de Goiás: o natural e o social na contemporaneidade. Goiânia: IESA, 2002. p. 91-118.

CARDOSO, M. R. D.; MARCUZZO, F. F. N.; BARROS, J. R. Classificação climática de Köppen-Geiger para o estado de Goiás e o Distrito Federal. Acta Geográfica (UFRR), v.8, p.40-55, 2014.

CASSOL, P. B.; BOHNER, T. Os eventos climáticos e a sua indissociabilidade na saúde e na economia global. Revista Eletrônica em Gestão, Educação e Tecnologia Ambiental, v. 5, n. 5, p. 653-657, 2012.

CASTELHANO, F. J.; PINTO, J. E. S. S. Tendências e alterações climáticas no Estado de Sergipe, nordeste do Brasil. Revista do Departamento de Geografia, v. 42, p. e185565-e185565, 2022. https://doi.org/10.11606/eISSN.2236-2878.rdg.2022.185565.

COELHO, F. J. S. et al. Seasonal study on changing trends in climatic extremes indicators in the Brazilian territory. Revista Brasileira de Cartografia, v. 75, 2023. http://dx.doi.org/10.14393/rbcv75n0a-68545

COSTA, R. L. et al. Tendências em índices extremos de precipitação em Cabaceiras (PB) para períodos distintos. Revista Brasileira de Geografia Física, v. 13, n. 1, p. 271-285, 2020. https://doi.org/10.26848/rbgf.v13.1.p271-285 .

COPERNICUS Global Climate Highlights 2024. 2025. Disponible: https://climate.copernicus.eu/global-climate-highlights-2024?utm_source=socialmedia&utm_medium=bs&utm_campaign=global-climate-highlights-2024

DEUS, R. M. M. de.; NASCIMENTO, D. T. F. Como a mídia tem noticiado os eventos climáticos extremos no estado Goiás. In: ENCONTRO NACIONAL DE GEÓGRAFOS, 2016, São Luís. Anais... , São Luís: AGB, 2016. p. 38-48.

DEUS, R. M. M.; NASCIMENTO, D. T. F. Eventos climáticos extremos no Estado de Goiás e Distrito Federal. In: SIMPÓSIO BRASILEIRO DE CLIMATOLOGIA GEOGRÁFICA, 14., 2021. João Pessoa. Anais... João Pessoa: UFPE, 2021. v. 1, p. 2.938-2.954.

FENNER, A.; SILVA, E.L.; GURGEL, H.C. Mudança climática e saúde: um perfil do Brasil. Organização Pan- Americana da saúde; Ministério da saúde. Brasília, 1. ed., 2009.

HOFF, H. Understanding the Nexus. Background Paper for the Bonn 2011 Conference: The Water, Energy and Food Security Nexus. Stockholm: Stockholm Environment Institute, 2011.

HOFMANN, G. S. et al. The Brazilian Cerrado is becoming hotter and drier. Global Change Biology, p. 1-14, 2021. https://doi.org/10.1111/gcb.15712.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

HOFMANN, G. S.; et al. Changes in atmospheric circulation and evapotranspiration are reducing rainfall in the Brazilian Cerrado. Sci Rep, 13, 11236 (2023). https://doi.org/10.1038/s41598-023-38174-X

IPCC. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V. et al. (eds.). Cambridge, UK and New York, NY, USA: Cambridge University Press, 2021.

JOSEPH, W.; SOUZA, A. P.; SABINO, M. Índices de extremos de temperatura do ar na Amazônia brasileira. Confins. Revue franco-brésilienne de géographie/Revista Franco-Brasileira de Geografia, n. 52, 2021. DOI: https://doi.org/10.4000/confins.41520.

KARL, T. R.; NICHOLLS, N.; GHAZI, A. CLIVAR/GCOS/WMO workshop on indices and indicators for climate extremes: Workshop summary. Climatic Change,v. 42, p. 3-7, 1999.

KOPPEN W.; GEIGER, R. (eds). Handbuck der Klimatologie. Berlin, 1961.

LUIZ, G. C. Influência da relação solo-atmosfera no comportamento hidromecânico de solos tropicais não saturados: estudo de caso - município de Goiânia-GO. 2012. 246 f. Tese (Doutorado em Geotecnia) - Universidade de Brasília, Brasília, DF, 2012.

LUIZ, G. C.; ROMÃO, P. de A. Interação Solo-Atmosfera e Processos de Inundação e Alagamento na Cidade de Goiânia-GO. Revista Brasileira De Geografia Física, v. 12, p. 1891-1903, 2019.

LUIZ-SILVA, W.; BAZZANELA, A.C.F.; DERECZYNSKI, C.P.; OSCAR-JÚNIOR, A.C.; RAUPP, I.P. Temperature and Precipitation Extremes in the Brazilian Legal Amazon: A Summary of Climatological Patterns and Detected Trends. Atmosphere, v. 16, n. 222, 2025.

MARCUZZO, F. F. N.; OLIVEIRA, N. de L.; CARDOSO, M. R. D. Tendência do número de dias de chuva no estado do Mato Grosso. Ciência e Natura, v. 32, n. 2, p. 59-82, 2013.

MARENGO, J. A. et al. Impactos de extremos relacionados com o tempo e o clima – Impactos sociais e econômicos. Boletim do Grupo de Pesquisa em Mudanças Climáticas. INPE, 2009.

MARENGO, J. A.; CARDONA, O. D.; MARTINEZ, R. Climatic hazards and disaster risk reduction in South-Central America and the Caribbean. Frontiers in Climate, v. 4, p. 1, 2022.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

MARIANI, L.; GUARENGHI, M. M.; MITO, J. Y. L.; CAVALIERO, C. K. N.; GALVÃO, R. R. de A. Análise de oportunidades e desafios para o nexo água-energia. Desenvolvimento e Meio ambiente, v. 37, edição especial, p. 9-30, 2016.

MASSELOT, P. et al. Excess mortality attributed to heat and cold: a health impact assessment study in 854 cities in Europe. Lancet Planet Health, v. 7, n. 4, p. 271-281, 2023.

MENDONÇA, F. Clima, tropicalidade e saúde: uma pesperctiva a partir da intensificação do aquecimento global. Revista brasileira de climatologia. v.1, n.1, p.100-112, 2005.

NASCIMENTO, D. T. F. Chuvas no estado de Goiás e no Distrito Federal a partir de estimativas por satélite e circulação atmosférica. 2016. 200 f. Tese (Doutorado em Geografia) - Universidade Federal de Goiás, Goiânia, 2016.

NASCIMENTO, D. T. F.; COSTA, Q. S. Permanência do regime de precipitação entre diversos anos-padrão habituais, secos e chuvosos: estudo de caso para Goiânia-GO. Geoambiente on-line, p. 1-14, 2020

NASCIMENTO, D. T. F.; OLIVEIRA, I. J. de. Os sistemas atmosféricos atuantes e responsáveis pela gênese das chuvas no estado de Goiás e no Distrito Federal. Revista Brasileira De Climatologia, v. 27, p. 747–771, 2021.

NASCIMENTO, D. T. F.; SANTOS, N. B. F. dos; BARROS, J. R. Evolução da temperatura do ar em Goiânia-GO (1961-2009). In: SIMPÓSIO INTERNACIONAL DE CLIMATOLOGIA, 2011, João Pessoa-PB. Anais do IV SIC, 2011.

MONTEIRO, C. A. de F. Notas para o estudo do clima do Centro-Oeste brasileiro. Revista Brasileira de Geografia. v. 13, n. 1, p. 3-46, jan-mar 1951.

NASCIMENTO, D. T. F.; LIMA, L. V. V.; CRUZ, V. P. Episódios e gênese dos eventos climáticos extremos em Goiânia-GO. Caderno de Geografia, v.29, n.57, p. 583-608, 2019. https://doi.org/10.5752/p.2318-2962.2019v29n57p583.

NASCIMENTO, D. T. F.; OLIVEIRA, I. J. de. Os sistemas atmosféricos atuantes e responsáveis pela gênese das chuvas no estado de Goiás e no Distrito Federal. Revista Brasileira de Climatologia, v. 27, n. 16, p. 747-771, 2020.

NEVES, G. Z. F. Distribuição geográfica das chuvas no Estado de Goiás e Distrito Federal: dinâmica e aplicações nos estudos rítmicos. 2018. Tese (Doutorado) – Universidade de São Paulo, São Carlos, 2018.

NEVES, G. Z. de F. D. A chuva no estado de Goiás e Distrito Federal: aspectos espaciais, temporais e dinâmicos. Revista Brasileira de Climatologia, Edição Especial, p. 67-82, 2019. http://dx.doi.org/10.5380/abclima.v0i0.66147

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

NEVES, G. Z. de F. D.; SANTOS, B. C. dos; BUARQUE, P. F. S. M.; SANCHES, R. G.; PERUSSI, R.; PEREIRA, D. N. B.; HORTA, I. T. L. G. Índices de extremos climáticos e tendências de precipitação no estado de Goiás e Distrito Federal (Brasil). Revista Brasileira de Climatologia, Dourados, MS, v. 34, Jan. / Jun. 2024.

NIMER, E. Climatologia do Brasil. Rio de Janeiro: IBGE, Departamento de Recursos Naturais e Estudos Ambientais. 1979.

PENEREIRO, J. C.; MESCHIATTI, M. C. Tendências em séries anuais de precipitação e temperaturas no Brasil. Engenharia Sanitária e Ambiental, v. 23, p. 319-331, 2018. https://doi.org/10.1590/S1413-41522018168763.

PETERSON, T. C. Climate Change Indices. WMO Bulletin, v. 54, n. 2, p. 83-86, 2005.

PETERSON, T. C.; et al. Report on the Activities of the Working Group on Climate Change Detection and Related Rapporteurs 1998-2001. Rep. WCDMP-47, WMO-TD 10712001. Geneve: WMO, 2001.

REBELLO, E. R. G. et al. Observação de mudanças de extremos climáticos (precipitação e temperatura) utilizando o software RClimDex. Estudo de caso: Distrito Federal. In: CONGRESSO BRASILEIRO DE METEOROLOGIA, XV., 2008. São Paulo: Anais... São Paulo: SBM. 2008, p. 1-6.

REGOTO, P. et al. Observed changes in air temperature and precipitation extremes over Brazil. International Journal of Climatology, v. 41, n. 11, p. 5.125-5.142, 2021. https://doi.org/10.1002/joc.7119.

RODRIGUES, J. C. M. O nexo água-energia-alimentos aplicado ao contexto da Amazônia Paraense. 2017. 92 f. Dissertação (Mestrado) - Universidade Federal do Pará, Instituto de Filosofia e Ciências Humanas, Belém, 2017.

ROMANELLO, M. et al. The 2024 report of the Lancet Countdown on health and climate change: facing record-breaking threats from delayed action. The Lancet, v. 404, n. 10465, p. 1847-1896, 2024.

SALVIANO, M. F.; GROPPO, J. D.; PELLEGRINO, G. Q. Análise de tendências em dados de precipitação e temperatura no Brasil. Revista Brasileira de Meteorologia, v. 31, p. 64-73, 2016. https://doi.org/10.1590/0102-778620150003.

SANTOS, A. L. M.; GONÇALVES, W.A.; ANDRADE, L.M.B.; RODRIGUES, D.T.; BATISTA, F.F.; LIMA, G.C.; SILVA, C.M.S. Space-time characterization ox extreme precipitation indices for the semiarid region of Brazil. Climate, v. 12, n. 43, 2024

SANTOS, C. A. C. dos; BRITO, J. I. B. de; RAO, T. V. R.; MENEZES, H. E. A. Tendências dos índices de precipitação no Estado do Ceará. Revista Brasileira de Meteorologia, v. 24, n. 1, p. 39-47, 2009.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

SANTOS, L. A. R. dos; et al.. Caracterização de extremos climáticos utilizando o software Rclimdex. Estudo de caso: sudeste de Goiás. In: XIV CONGRESSO BRASILEIRO DE METEOROLOGIA, 2006, FLORIANÓPOLIS - SC. Anais...., Florianópolis: CBM, 2006.

SANTOS, L. O. F.; et al. Trends in Precipitation and Air Temperature Extremes and Their Relationship with Sea Surface Temperature in the Brazilian Midwest. Atmosphere, 14, 426, 2023. https://doi.org/10.3390/atmos14030426.

SERRA, A.; RATISBONNA, L. As massas de ar na América do Sul. Rio de Janeiro: Serviço de Meteorologia, Ministério da Agricultura, 1942.

SILVA, C. B.; CARPENEDO, C. B. Teleconexões no Cerrado brasileiro. IN: NASCIMENTO, D. T. F.; MARTINS, A. P.; LUIZ, G. C.; LOPES, R. M. Climatologia do Cerrado: variabilidades, suscetibilidades e mudanças climáticos no contexto do Cerrado brasileiro. Goiânia: C&A Alfa Comunicação, 2021. Cap. 9, p. 232-262.

SILVA, G. C.; NASCIMENTO, D. T. F. Variabilidade e tendência climática em Goiânia-GO: avaliação a partir de índices de extremos climáticos calculados em RClimdex. In: SIMPÓSIO BRASILEIRO DE CLIMATOLOGIA GEOGRÁFICA, 14., 2021, João Pessoa. Anais... João Pessoa: UFPE, 2021. v. 1, p. 2.922-2.937.

SILVA, P. E.; SANTOS E SILVA, C.M.; SPYRIDES, M.H.C.; ANDRADE, L.M.B. Precipitation and air temperature extremes in the Amazon and northeast Brazil. Int. J. Climatol., v. 39, p. 579–595, 2019.

SILVA, R. O. B.; MONTENEGRO, S. M. G. L.; SOUZA, W. M. Tendências de mudanças climáticas na precipitação pluviométrica nas bacias hidrográficas do estado de Pernambuco. Revista de Engenharia Sanitária e Ambiental, v.22, n.3, p. 579-589, 2017.

SILVA, W. L.; et al. Tendências observadas em indicadores de extremos climáticos de temperatura e precipitação no estado do Paraná. Revista Brasileira de Meteorologia, v.30, n.2, 181 - 194, 2015.

SKANSI, M. M. et al. Warming and wetting signals emerging from analysis of changes in climate extreme indices over South America. Global and Planetary Change, v. 100, p. 295-307, 2013. http://dx.doi.org/10.1016/j.gloplacha.2012.11.004

WMO - World Meteorological Organization. State of the Global Climate 2023. WMO: Geneva, 2024. n. 1347, 2024. Disponível em: https://library.wmo.int/idurl/4/68835

WMO - World Meteorological Organization. WMO Guidelines on the Calculation of Climate Normals. Geneva: WMO, 2017.

SciELO Preprints - This document is a preprint and its current status is available at: https://doi.org/10.1590/SciELOPreprints.16770

ZHANG, X.; YANG, F. RClimDex (1.0). User Guide. Ontario: Scientific Research, 2004.

ZHANG, X. et al. Avoiding inhomogeneity in percentile-based indices of temperature extremes. Journal of Climate, v. 18, n. 11, p. 1.641-1.651, 2005. https://doi.org/10.1175/JCLI3366.1.

This preprint was submitted under the following conditions:

- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a Creative Commons CC-BY license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.