

Publication status: Preprint has been published in a journal as an article
DOI of the published article: <https://doi.org/10.1590/2236-8906-22/2022>

Periphytic algal flora of the low Doce river, Espírito Santo State, Brazil, after ore tailings flow, 1: smooth cell walled *Cosmarium* Corda (Zygnematophyceae, Desmidiaceae)

Bruna Fadul-Souza, Karoline Barros de Souza, Mateus Sardi Trancoso, Stéfano Zorzal-Almeida, Carlos Eduardo de Mattos Bicudo, Geraldo José Peixoto Ramos, Valéria de Oliveira Fernandes

<https://doi.org/10.1590/2236-8906-22/2022>

Submitted on: 2022-08-19

Posted on: 2022-08-19 (version 1)
(YYYY-MM-DD)

Periphytic algal flora of the Low Doce River, Espírito Santo State, Brazil, after ore tailings flow, 1: smooth cell walled *Cosmarium* Corda (Zygnematophyceae, Desmidiaceae)¹

Bruna Fadul-Souza^{1,2,6}, Karoline Barros de Souza², Mateus Sardi Trancoso², Stéfano Zorzal-Almeida³, Carlos Eduardo de Mattos Bicudo⁴, Geraldo José Peixoto Ramos⁵ and Valéria de Oliveira Fernandes²

Short title: Smooth cell walled *Cosmarium* of the Low Doce River

Bruna Fadul-Souza: <https://orcid.org/0000-0003-2115-1687>

Carlos Eduardo de Mattos Bicudo: <https://orcid.org/0000-0003-4030-9961>

Geraldo José Peixoto Ramos: <https://orcid.org/0000-0003-2089-0461>

Karoline Barros de Souza: <https://orcid.org/0000-0003-4062-5382>

Mateus Sardi Trancoso: <https://orcid.org/0000-0002-8890-6266>

Stéfano Zorzal-Almeida: <https://orcid.org/0000-0002-9977-5898>

Valéria de Oliveira Fernandes: <https://orcid.org/0000-0002-1968-3927>

¹ First Author's course conclusion work

² Universidade Federal do Espírito Santo, Departamento de Ciências Biológicas, Laboratório de Taxonomia e Ecologia de Algas Continentais, Avenida Fernando Ferrari, 514, 29075-910 Vitória, ES, Brazil

³ Universidade Federal do Espírito Santo, Departamento de Oceanografia e Ecologia, Laboratório de Ecologia e Biodiversidade de Microprodutores Aquáticos, Avenida Fernando Ferrari, 514, 29075-910 Vitória, ES, Brazil

⁴ Instituto de Pesquisas Ambientais, Núcleo de Conservação da Biodiversidade, Avenida Miguel Estéfano, 3687, 04301-902 São Paulo, SP, Brazil

⁵ Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Rodovia Jorge Amado, Km 16, 45662-900 Ilhéus, BA, Brazil

⁶ Corresponding author: b.fadul.souza@gmail.com

ABSTRACT – (Periphytic algal flora of the low Doce river, Espírito Santo State, Brazil, after ore tailings flow, 1: smooth cell walled *Cosmarium* Corda (Zygnematophyceae, Desmidiaceae). Study aimed at identifying and recording smooth or finely punctate cell walled *Cosmarium* Corda species of the low Doce river region, Espírito Santo State, Brazil. Periphytic material was collected monthly during the period October 2018 to February 2020, except for October 2019. We identified 22 taxa and analyzed the environmental conditions (trophic state, pH and total iron) in which each taxon occurred. Our study recorded 21 new records of smooth cell walled *Cosmarium* for Espírito Santo State and one (*Cosmarium trilobulatum* var. *tumidum* Willi Krieger & Gerloff) for Brazil.

Keywords: desmids, periphyton, taxonomy

RESUMO – (Ficoflórula perifítica do Baixo Rio Doce, Estado do Espírito Santo, Brasil, após passagem de rejeitos de minério, 1: *Cosmarium* Corda de parede lisa (Zygnematophyceae, Desmidiaceae). Este estudo teve como objetivo identificar e registrar as espécies de *Cosmarium* Corda de parede lisa ou finamente pontuada na região do baixo rio Doce, no Estado do Espírito Santo, Brasil. O material perifítico foi coletado mensalmente durante o período outubro de 2018 a fevereiro de 2020, exceto no mês de outubro de 2019. Identificamos 22 táxons e analisamos as condições ambientais (estado trófico, pH e ferro total) nas quais cada táxon ocorreu. O estudo registrou 21 novos registros de *Cosmarium* de parede lisa para o Estado do Espírito Santo e um (*Cosmarium trilobulatum* var. *tumidum* Willi Krieger & Gerloff) para o Brasil.

Palavras-chave: desmídias, perifíton, taxonomia

Introduction

Originally described by Ralfs in 1848, genus *Cosmarium* Corda gathers over 1,500 taxa and consists in the most numerous and one of the oldest genera of family Desmidiaceae. Genus is composed by usually free-living single cells with a median constriction that vary from a deep closed depression to a widely open and shallow one, that divides the cell into two symmetrical semicells and vary in shape and size depending on the species (Bicudo & Menezes 2017). Taxonomy of *Cosmarium* is primarily based on the cell outline and cell wall ornamentation. For practical purposes, however, taxonomists tend to divide it into two artificial groups: one with ornamented cell walls (ornamentation varying from small granules to scrobiculations and tiny spines) and the other one with smooth or finely punctate cell wall.

Genus *Cosmarium* is worldwide recorded in many taxonomic surveys. From classical 19th century literature (e.g., Brébisson 1856, Lundell 1871, Nordstedt & Wittrock 1876, Ralfs 1848, West & G.S. West 1895) to the more recent papers, the genus was recorded for many countries, including Nigeria (Biswas & Nweze 1990), Colombia (Duque & Donato 1996), Canada (Elster *et al.* 1997), Bangladesh (Islam & Irfanullah 2006), India (Dhande & Jawale 2009, Bansod & Patil 2019), Czech Republic (Šťastný 2009), South Korea (Lee 2015) and many others. Over 600 *Cosmarium* taxa were recorded for Brazil (Biolo & Bicudo 2018), and the genus was more intensively studied in the south region where numerous taxonomic studies revealed its presence in both phytoplankton and periphyton of reservoirs and rivers. Recent floristic studies recording the presence of *Cosmarium* in Brazil include States such as Paraná (Silva & Cecy 2004, Bortolini *et al.* 2010, Felisberto & Rodrigues 2010, Biolo *et al.* 2013, Menezes *et al.* 2011, Carvalho *et al.* 2015, Moresco *et al.* 2015, Aquino *et al.* 2016, Osório *et al.* 2018, Rodrigues & Moresco 2021), São Paulo (Bicudo *et al.* 2019, Felisberto & Rodrigues 2010), Bahia (Oliveira *et al.* 2010, Oliveira *et al.* 2011, Oliveira *et al.* 2016, Ramos *et al.* 2018, Ramos *et al.* 2021), Goiás (Felisberto & Rodrigues 2004), Distrito Federal (Estrela *et al.* 2011), Amazonas (Saturnino *et al.* 2020), Sergipe (Burlle *et al.* 2020), Mato Grosso and Rio de Janeiro (Kanetsuna 2004). Furthermore, in the last 10 years five new species of *Cosmarium* were described based on Brazilian material: *Cosmarium bahianum* G.J.P.Ramos, C.E.M.Bicudo & C.W.N.Moura, *Cosmarium bromelicola* G.J.P.Ramos, C.E.M.Bicudo & C.W.N.Moura, *Cosmarium diamantinum* G.J.P.Ramos & C.W.N.Moura, *Cosmarium imperiale* G.J.P.Ramos & C.W.N.Moura and *Cosmarium oliveirae* G.J.P.Ramos, C.E.M.Bicudo & C.W.N.Moura, in addition to a few taxonomic varieties. Despite having one of the most important coastal lake systems of Brazil (Barroso *et al.* 2012), Espírito Santo State has been widely neglected concerning its algal flora studies. Delazari-Barroso *et al.* (2007) is the only floristic study of the State recording the presence of the genus by four *Cosmarium* taxa (*Cosmarium asphaerosporum* var. *strigosum* Nordstedt, *Cosmarium* cf. *capitulum* J.Roy & Bisset, *Cosmarium pseudobroomei* Wolle and *Cosmarium pseudoconnatum* Nordstedt).

The Low Doce River region is undergoing severe urban and rural anthropogenic interference mainly represented by livestock, agriculture and the increasing industrialization (Barroso *et al.* 2012) even before the Fundão dam failure in 2015. Dam rupture released downstream about 43 million m³ (Carmo *et al.* 2017) of mining tailings, causing several changes on the fluvial system, such as erosion, degradation, and iron (Fe) enrichment to potentially impactful levels (Hatje *et al.* 2017). Lack of floristic studies in the region and the fact that mine tailing inputs have previously caused taxonomic and functional changes in several communities, such as the benthic fauna (Burd 2002), marine macroalgal flora (Correa *et al.* 1999), planktonic algae (Salonen *et al.* 2006), and benthic algal

communities (Lavoie *et al.* 2012), highlights the importance of periphyton biodiversity studies in the region.

Present study aimed at performing a taxonomic survey of the smooth cell walled *Cosmarium* species of the Low Doce River region three years after the input of iron ore tailings, also focusing on species autecology, thus contributing to the knowledge of freshwater algal diversity of Espírito Santo State.

Materials and Methods

The study was carried out in three cities in Espírito Santo State: Baixo Guandu, Colatina and Linhares. Detailed information on study area and sampling design are available in Zorzal-Almeida *et al.* (2021). For taxonomic and ecological analysis, a minimum of five non-permanent slides were prepared for each sample. Material was observed and photographed, respectively, with a Motic Panthera L digital binocular upright Olympus microscope and a Moticom 3.0 MP camera. The following taxonomic features were observed for identification and description of the material at the microscope: cell shape, number and position of chloroplasts and pyrenoids, cell wall structure, and metric data such as cell length and width, and isthmus width. Cell length (L), cell width (W) and isthmus width (I) were measured using Motic Image Plus 3.0 software. Identification was based on classic literature (e.g. Ralfs 1848, West & West 1895, 1902, Croasdale 1956, Scott & Grönblad 1957, Krieger 1932) and recent papers (e.g. Coesel & Meesters 2007, Oliveira *et al.* 2016, Bicudo *et al.* 2019, etc.).

Environmental conditions were described according to ecosystem type (rivers, lagoons and lakes), trophic state index (TSI), and pH (method specifications in Zorzal-Almeida *et al.* 2021), classified as acid (< 6.5), alkaline (> 7.5) and circumneutral (6.5-7.5) (Šťastný 2010). Occurrence frequency of each taxon was calculated according to Zorzal-Almeida *et al.* (2021). To highlight the first citations for Espírito Santo State, taxa are marked with [*] and the first citations for Brazil with [**]. Samples studied during the present inventory are deposited at the Herbarium VIES/UFES.

Results and discussion

Twenty-two smooth cell walled *Cosmarium* Corda taxa were presently identified, from which 21 were recorded for the first time for Espírito Santo State and one for Brazil. List of taxa and their occurrence frequency are listed in Table1.

* *Cosmarium abbreviatum* Raciborski var. *minus* (West & G.S.West) Willi Krieger & Gerloff, Gattung *Cosmarium* 3-4: 242. 1969; 2: pl. 42, fig. 18. 1965.

Figure 1

Cell 1.1-1.2 times longer than broad, sometimes c. 1.1 times broader than long; median constriction deep, sinus linear, slightly open at the extremity; semicells trapezoidal, apex truncate, angles rounded, lateral margins convex to slightly straight; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell; cell apical view elliptical, semicell lateral view circular. L: 6.8-14.1 µm; W: 6.3-11.3 µm; I: 1-4.3 µm.

Note: This variety showed a considerable morphological plasticity, mainly in their length:width ratio and lateral margins shape. Bicudo (1988) discussed the polymorphism in *Cosmarium abbreviatum* var. *minus* and its consequences for the taxonomic identification of such small desmids, suggesting the great probability of many taxa, among which *Cosmarium succisum* G.S.West var. *jaoi* Willi Krieger & Gerloff and *C. abbreviatum* var. *minus*, might just be morphological expressions of the one same species. Present material displayed the same morphological variation described in the above-cited literature.

Present material occurred in all lentic and lotic environments studied, under oligotrophic to hypertrophic, and acidic to alkaline conditions.

Material examined: BRDE00A01, BRDE21A03, BRDE22A03, BRDE20A04, BRDE23A03, BRDE00A05, BRDE21A05, BRDE26A05, BRDE23A04, BRDE25A09, BRDE23A05, BRDE00A07, BRDE26A07, BRDE23A06, BRDE20A08, BRDE21A09, BRDE22A08, BRDE23A08, BRDE25A17, BRDE00A10, BRDE20A10, BRDE20A12, BRDE25A24, BRDE00A13, BRDE22A12, BRDE24A14, BRDE25A29, BRDE20A16, BRDE25A32, BRDE21A01, BRDE18A01, BRDE19A01, BRDE18A02, BRDE18A03, BRDE18A04, BRDE19A04, BRDE18A05, BRDE19A07, BRDE24A08, BRDE18A10, BRDE25A20, BRDE18A11, BRDE19A11, BRDE24A10, BRDE18A12, BRDE19A12, BRDE24A11, BRDE17A13, BRDE18A13, BRDE25A26, BRDE22A13, BRDE18A14, BRDE25A28, BRDE22A14, BRDE19A15.

* *Cosmarium angulosum* Brébisson, Mémoires de la Société Impériale des Sciences Naturelles de Cherbourg 4: 127. pl. 1, fig. 17. 1856.

Figure 2

Cell c. 1.4 times longer than broad, median constriction deep, sinus linear; semicells quadratic, apex retuse to slight truncate, lateral margins parallel, apical angles concave, basal angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: c. 22.8 µm; W: c. 15.8 µm; I: c. 5.1 µm.

Note: Only a single specimen of this species was found during the present study, which was a little different from the ones in Brébisson (1856) and Coesel & Meesters (2007). Present specimens were slightly convex at the middle of the apex instead of a truncate one, and concave instead of straight apical angles.

Present material was collected from a lentic environment, under oligotrophic alkaline conditions.

Material examined: BRDE25A29.

* *Cosmarium baileyi* Wolle, Smithsonian Contributions to Knowledge 2(8): 36, pl. 1, fig. 1. 1851.

Figure 3

Cell c. 1.1 times broader than long, median constriction deep, sinus linear; semicells semicircular, apex truncate, angles rounded; cell wall punctate, chloroplasts 2, pyrenoids 2 per semicell. L: c. 29.9 µm; W: c. 32.8 µm; I: c. 10.4 µm.

Note: *Cosmarium baileyi* is morphologically similar to *Cosmarium obsoletum* (Hantzsch) Reinsch, but the latter has thickened, angular, convergent basal angles, whereas the first one has rounded ones. Present material was gathered in a lentic environment under mesotrophic, eutrophic, hypertrophic, and circumneutral conditions.

Material examined: BRDE19A11, BRDE19A14, BRDE25A28.

* *Cosmarium clepsydra* Nordstedt var. *dissimile* (Raciborski) Willi Krieger & Gerloff, Gattung *Cosmarium* 2: 144, pl. 30, fig. 4. 1965.

Figures 4-6

Cell c. 1.1 times longer than broad, median constriction deep, sinus linear; semicells semicircular, apex rounded, angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 11.1-17.8 µm; W: 9.7-15.2 µm; I: 3.1-5.4 µm.

Note: Present variety differs from the typical by having semicircular semicells instead of triangular or almost so.

Present material was gathered from a lentic environment under oligotrophic to eutrophic and alkaline conditions.

Material examined: BRDE18A06, BRDE18A13, BRDE18A16.

* *Cosmarium contractum* Kirchner var. *minutum* (Delponte) Coesel, Cryptogamie, Algologie 10: 184, pl. 2, fig. 2. 1989.

Figures 7, 8

Cell 1.4-1.5 times longer than broad, median constriction deep, sinus widely open; semicells ellipsoid, apex rounded, angles rounded; cell wall punctate, chloroplast 1, pyrenoid 1 per semicell; frequently involved in some spherical, inconspicuous mucilaginous envelope. L: 15.9-24.7 μm ; W: 10.9-17.4 μm ; I: 3-4.4 μm .

Note: This variety differs from the typical in having smaller cell dimensions, length (12-)29-30 μm and breadth (9-)18-25 μm , according to Coesel & Meesters (2007).

Present material was collected from deep lakes, under ultraoligotrophic to eutrophic and acidic to alkaline conditions.

Material examined: BRDE18A01, BRDE19A01, BRDE18A02, BRDE19A02, BRDE18A03, BRDE19A03, BRDE18A04, BRDE19A04, BRDE19A05, BRDE20A05, BRDE19A06, BRDE19A07, BRDE19A09, BRDE18A11, BRDE19A11, BRDE19A12, BRDE20A13, BRDE19A14, BRDE18A15, BRDE19A15, BRDE18A16, BRDE19A16.

* *Cosmarium contractum* Kirchner var. *sparsipunctatum* Kurt Förster *ex* Kurt Förster, Amazoniana 2(1-2): 48, pl. 13, fig. 13-14. 1969.

Figures 9, 10

Cell 1.2-1.3 times longer than broad, median constriction deep, sinus widely open; semicells elliptical, apex truncate to slight rounded, angles rounded; cell wall punctate, chloroplast 1, pyrenoid 1 per semicell. L: 12-16.9 μm ; W: 9.4-14 μm ; I: 2.8-3.9 μm .

Note: According to Oliveira *et al.* (2016), *Cosmarium contractum* var. *sparsipunctatum* differs from the typical variety in its smaller dimensions, punctate cell wall, elliptical semicells, and a more round and broad isthmus.

Present material was collected living in somewhat deep lakes (Nova and Limão), under ultraoligotrophic, eutrophic and circumneutral conditions.

Material examined: BRDE18A08, BRDE19A16.

* *Cosmarium difficile* Lütkemüller, Verhandlungen der zoologisch-botanischen Gesellschaft in Wien 42: 551, pl. 8, fig. 3. 1892.

Figures 11, 12

Cell c. 1.8 times longer than broad, median constriction deep, sinus closed near the isthmus, open at the extremity; semicells quadratic, apical and basal angles rounded, apex truncate, a refractive thickening in the center; cell wall smooth, with 3 horizontal rows of punctations on each semicell, chloroplast 1, pyrenoid 1 per semicell. L: 19.4-30.7 μm ; W: 10.8-17.9 μm ; I: 3.5-3.7 μm .

Note: The present species has a feature not usually observed in *Cosmarium difficile* Lütkemüller, *i.e.*, a refractive thickening at the apex of the semicells. The only representative of *C. difficile* having such apical thickening is *C. difficile* f. *laeve* Scott & Prescott, however, this forma does not have the typical pore rows of the species. Thus, it was decided that the current material would be best identified as *C. difficile*, mainly due to the typical punctuation rows despite of the originally non-described feature. The possibility of the present material be representative of a new variety of *C. difficile* is also an open question to be considered in the future, until further studies be made towards its best delimitation at intraspecific level.

Present material was collected from lentic systems, under oligotrophic to eutrophic and circumneutral to alkaline conditions.

Material examined: BRDE18A02, BRDE24A02, BRDE25A06, BRDE18A06, BRDE19A06, BRDE19A07, BRDE24A06, BRDE19A11, BRDE18A15, BRDE19A15, BRDE25A29, BRDE18A16.

* *Cosmarium galeritum* Nordstedt var. *subtumidum* Borge, Arkiv für Botanik 1: 95, pl. 3, fig. 14. 1903.

Figures 13, 14

Cell c. 1.1 times longer than broad or as long as broad, median constriction deep, sinus linear; semicells trapezoidal, apex truncate, angles rounded; cell wall punctate, chloroplasts 2, pyrenoids 2 per semicell. L: 31.6-45.5 μm ; W: 31.2-43 μm ; I: 10.8-14.3 μm .

Note: *Cosmarium galeritum* var. *subtumidum* differs from the nominate variety in having more rounded semicells (Borge 1903). Present specimen showed rounded to slightly truncate apex, a variation similar to that in Borge's (1903) illustration, a fact that led us to its current identification.

Present material was found in lentic systems, under ultraoligotrophic to eutrophic and circumneutral conditions.

Material examined: BRDE25A02, BRDE18A05, BRDE19A07, BRDE18A11, BRDE18A13, BRDE20A13, BRDE25A26, BRDE19A15, BRDE25A29, BRDE19A16, BRDE18A13.

* *Cosmarium granatum* Brébisson ex Ralfs var. *granatum*, British Desmidiaceae. 56, pl. 32, fig. 6. 1848.

Figures 15, 16

Cell 1.4-1.5 times longer than broad, median constriction deep, sinus linear; semicells pyramidal, apex sharp or truncate round, basal angles rounded to straight; cell wall smooth, sometimes thickened at the apex, chloroplast 1, pyrenoid 1 per semicell. L: 20-38.2 µm; W: 13-26.7 µm; I: 4.4-9.7 µm.

Note: Specimens studied varied in several morphological features. Cell shape varied from pyramidal to slight trapezoid, basal angles from sometimes wide concave to straight, and the apex from rounded to truncate to slightly sharp. According to Bortolini *et al.* (2010), this species displays some great morphological variation, a fact that was also observed during the present study.

Present material occurred in lentic and lotic environments, under ultraoligotrophic to supertrophic and acidic to alkaline conditions.

Material examined: BRDE21A01, BRDE18A01, BRDE00A02, BRDE21A02, BRDE22A01, BRDE18A02, BRDE00A03, BRDE21A03, BRDE22A02, BRDE26A03, BRDE17A04, BRDE18A04, BRDE22A04, BRDE18A05, BRDE00A06, BRDE21A06, BRDE22A05, BRDE19A06, BRDE00A07, BRDE21A07, BRDE22A06, BRDE26A07, BRDE18A07, BRDE17A08, BRDE00A08, BRDE21A08, BRDE22A07, BRDE26A08, BRDE18A08, BRDE17A09, BRDE00A09, BRDE21A09, BRDE22A08, BRDE18A09, BRDE00A10, BRDE21A10, BRDE22A09, BRDE26A10, BRDE18A10, BRDE00A11, BRDE21A11, BRDE22A10, BRDE19A11, BRDE22A11, BRDE18A12, BRDE25A24, BRDE21A13, BRDE22A12, BRDE26A13, BRDE18A13, BRDE26A14, BRDE18A14, BRDE00A15, BRDE21A15, BRDE22A14, BRDE26A15, BRDE18A15, BRDE17A16, BRDE22A15, BRDE18A16, BRDE18A01, BRDE26A12, BRDE17A13, BRDE26A15.

* *Cosmarium impressulum* Elfving, Acta Societatis Fauna Flora Fennica 2(2): 13, pl. 1, fig. 9. 1881.

Figure 17

Cell c. 1.5 times longer than broad, median constriction deep, sinus linear; semicells semicircular, margins undulate, 6-7 undulations per semicell, basal angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: c. 17 µm; W: c. 11.4 µm; I: c. 3.2 µm.

Note: Just a single specimen of this species was detected during the present study. When compared to Silva & Cecy's (2004) population, current specimen is much longer and has less prominent marginal undulations. It is also smaller when compared to Felisberto & Rodrigues (2010) material, and has fewer undulations at the lateral margins when compared to Menezes *et al.* (2011). *Cosmarium impressulum* may be mistaken by *Cosmarium undulatum* Corda ex Ralfs, but according to Bicudo *et*

al. (2019), however, they are distinct in the number of marginal undulations (*C. undulatum* has 10-12 undulations, whereas *C. impressulum* has up to 6), and their cell dimensions (*C. undulatum* is comparatively larger: 44-64 x 30-52 µm).

Present material occurred in Doce River under mesotrophic and alkaline conditions.

Material examined: BRDE00A11.

* *Cosmarium laeve* Rabenhorst var. *laeve*, Flora Europaea Algarum 3: 161. 1868.

Figures 18, 19

Cell 1.5-1.6 times longer than broad, median constriction deep, sinus linear; semicells semicircular to subpyramidal, apical margin often asymmetrical, apex rounded to slightly truncate, basal angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell; lateral view subcircular. L: 18-20.4 µm; W: 12.9-13.8 µm; I: 3.7-4.4 µm.

Note: Present material was collected both from lotic and lentic environments, under ultraoligotrophic-supertrophic and circumneutral-alkaline conditions.

Material examined: BRDE17A01, BRDE25A01, BRDE00A02, BRDE21A02, BRDE22A01, BRDE17A03, BRDE25A05, BRDE17A04, BRDE22A03, BRDE17A05, BRDE00A05, BRDE21A05, BRDE22A04, BRDE26A05, BRDE17A06, BRDE00A06, BRDE21A06, BRDE22A05, BRDE25A11, BRDE17A07, BRDE00A07, BRDE21A07, BRDE22A06, BRDE26A07, BRDE25A13, BRDE17A08, BRDE21A08, BRDE25A15, BRDE17A09, BRDE00A09, BRDE21A09, BRDE26A09, BRDE17A10, BRDE17A11, BRDE19A11, BRDE17A12, BRDE17A13, BRDE21A13, BRDE26A13, BRDE00A14, BRDE00A15, BRDE21A15, BRDE22A14, BRDE26A15, BRDE25A29, BRDE17A16.

* *Cosmarium lundellii* Delponte var. *madagascariense* West & G.S. West, Transactions of the Linnean Society of Botany 5: 55, pl. 8, fig. 2. 1895.

Figures 20, 21

Cell c. 1.2 times longer than broad, median constriction deep, sinus linear, gradually opening towards the extremity; semicells pyramidal, apex rounded to slightly truncate, basal angles rounded; cell wall punctate, chloroplasts 2, pyrenoids 2 per semicell. L: 50-61.8 µm; W: 40.2-51.6 µm; I: 13-20.4 µm.

Note: Present variety differs from the typical by being longer than broad not as long as broad. Specimens studied were smaller than the one described in West & West (1895) (L: c. 81 µm; W: c. 70 µm; I: 33 µm).

Present material was collected from a lentic system, under mesotrophic and alkaline conditions.

Material examined: BRDE18A13.

* *Cosmarium moniliforme* (Turpin) Ralfs, British Desmidiaceae. 107, pl. 17, fig. 6. 1848.

Figure 22

Cell 1.6-1.7 times longer than broad, median constriction deep, sinus wide open; semicell subcircular, apex wide rounded, lateral margins convex, angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 29.7-34.8 μm ; W: 17.5-21.3 μm ; I: 4.5-6.5 μm .

Note: According to Biolo *et al.* (2013), *Cosmarium moniliforme* is very similar to *Cosmarium contractum* Kirchner, but differs in having comparatively more subcircular semicells and elliptical apical view. The apical view of the present specimens was not observed, but the subcircular semicells outline led to its identification as *C. moniliforme*.

Present material was gathered from both lotic and lentic systems, always under oligotrophic, mesotrophic and alkaline conditions.

Material examined: BRDE21A07, BRDE25A29, BRDE25A32.

* *Cosmarium norimbergense* Reinsch var. *depressum* (West & G.S.West) Willi Krieger & Gerloff, Gattung *Cosmarium* 3-4: 292, pl. 48, fig. 1. 1969.

Figure 23

Cell c. 1.1 times longer than broad, median constriction deep, sinus linear, slightly open near the extremity; semicells rectangular, apex truncate, lateral margins concave, apical and basal angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 11.1-12.9 μm ; W: 9.7-11.6 μm ; I: 3.5-4.8 μm .

Note: *Cosmarium norimbergense* var. *depressum* differs from the type of the species in its vertically flattened cells, usually 1.1-1.2 times longer than broad (Aquino *et al.* 2014).

Present material was collected from a lentic system, under ultraoligotrophic, oligotrophic, and circumneutral-alkaline conditions.

Material examined: BRDE25A30, BRDE26A16, BRDE18A16, BRDE24A15.

* *Cosmarium ocellatum* Eichler & Gutwinski, Rozprawy Wydziału matematyczno-przyrodniczego akademii umiejetnosci 28: 164, pl. 4, fig. 7. 1894.

Figure 24

Cell c. 1.4 times longer than broad, median constriction deep, sinus wide open; semicells trapezoid, apex truncate, apical angles slightly convex, basal angles rounded; cell wall punctate, chloroplast 1, pyrenoid 1 per semicell. L: c. 26.2 μm ; W: c. 18.8 μm ; I: c. 5.0 μm .

Note: Only one individual specimen of *Cosmarium ocellatum* was found during this survey.

Present material was gathered from a lentic system under mesotrophic circumneutral conditions.

Material examined: BRDE18A05.

* *Cosmarium pachydermum* P.Lundell var. *aethiopicum* (West & G.S.West) West & G.S.West, A monograph of the British Desmidiaceae 2: 140, pl. 57, fig. 8, 9. 1905.

Figure 25

Cell 1.2-1.4 times longer than broad, median constriction moderate, sinus closed near the isthmus, gradually open towards the extremity; semicells semicircular, apex rounded to slightly truncate, basal angles rounded; cell wall punctate, chloroplasts 2 per semicell, pyrenoid not seen. L: 86-130 μm ; W: 65-94 μm ; I: 31-39 μm .

Note: This variety differs from the typical by having thinner cell wall, a more open sinus, and the relatively broader cell (Krieger & Gerloff 1962). Specimens in the population currently studied showed a wider range of its length:width ratio, whereas according to Krieger & Gerloff (1962) that of the present variety is about 1.25 times longer than broad.

Present material was gathered both from lentic and lotic systems, in oligotrophic to mesotrophic and alkaline conditions.

Material examined: BRDE18A02, BRDE18A03, BRDE00A11, BRDE19A16.

* *Cosmarium polygonum* (Nägeli) W.Archer var. *depressum* Messikommer, Beiträge zur geobotanischen Landesaufnahme der Schweiz 24: 147, pl. 5, fig. 8. 1942.

Figures 26, 27

Cell 1.1-1.3 times broader than long, median constriction deep, sinus closed near the isthmus, gradually open towards the extremity; semicells hexagonal, apex truncate, apical angles acute; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell; papillae in the mid region of semicells, apical view diamond-shaped, lateral view not observed. L: 8-9.2 μm ; W: 10-11 μm ; I: 3.2-3.5 μm .

Note: Present variety differs from the typical by having half its size and being broader than long (Messikommer 1942). Material currently studied differs from the type in its slight greater dimensions (L: 8-8.7 μm ; W: 9.3-9.8 μm ; I: 3 μm).

Present material was collected from Areão lake, under mesotrophic and circumneutral to alkaline conditions.

Material examined: BRDE23A04, BRDE23A05.

Cosmarium pseudoconnatum Nordstedt, Videnskabelige Meddelelser fra den naturhistoriske Forening i Kjöbenhavn 1869(14-15): 214. 1869; 1887: pl. 3, fig. 17. 1870.

Figures 28, 29

Cell 1.3-1.4 times longer than broad, median constriction shallow, sinus wide open; semicells semicircular, apex rounded, angles rounded; cell wall densely punctate, isthmus area smooth, chloroplasts 4, pyrenoids 4 per semicell; apical view circular, lateral view the same as the front view. L: 30-63 µm; W: 23-47 µm; I: 21-44 µm.

Note: *Cosmarium pseudoconnatum* is very similar to *Cosmarium connatum* Brébisson ex Ralfs differing only in the presence of four chloroplasts in each semicell, whereas the latter has two. According to Scott & Grönblad (1957), cell wall of both species is identical: scrobiculate with easy visible pores in between scrobicules.

Present material occurred in both lentic and lotic systems, under oligotrophic-supertrophic and acidic-alkaline conditions. Delazari-Barroso *et al.* (2007) previously recorded it from the phytoplankton of a small water supply reservoir in Espírito Santo State.

Material examined: BRDE18A01, BRDE18A02, BRDE18A03, BRDE21A05, BRDE22A04, BRDE26A05, BRDE18A05, BRDE00A07, BRDE19A09, BRDE20A09, BRDE22A09, BRDE18A11, BRDE18A13, BRDE19A13, BRDE18A14, BRDE19A14, BRDE18A15, BRDE17A16, BRDE18A16, BRDE25A32, BRDE22A01, BRDE26A07, BRDE20A07, BRDE17A08, BRDE00A11, BRDE20A12.

* ***Cosmarium pseudopyramidatum*** P.Lundell, Nova Acta Regia Societatis Scientiarum Upsaliensis: sér. 3, 8(2): 41, pl. 2, fig. 18. 1871.

Figure 30

Cell c. 1.6 times longer than broad, median constriction deep, sinus linear; semicells pyramidal, apex rounded, apical and basal angles broadly rounded; cell wall punctate, chloroplast 1, pyrenoid 1 per semicell. L: c. 36 µm; W: c. 22.5 µm; I: c. 8.2 µm.

Note: According to Coesel & Meesters (2007), *Cosmarium pseudopyramidatum* is morphologically close to *Cosmarium pyramidatum* Brébisson ex Ralfs, but differs in having a single pyrenoid per semicell and being up to 60 µm long. Just one individual specimen of this species was collected during the present survey.

Present material was gathered in a lentic environment (Limão lake), under oligotrophic and alkaline conditions.

Material examined: BRDE18A16.

* *Cosmarium rectangulare* Grunow var. *hexagonum* (Delponte) West & G.S.West, A monograph of the British Desmidiaceae 3: 56, pl. 70, fig. 4. 1908.

Figures 31, 32

Cell 1.2-1.3 times longer than broad, median constriction deep, sinus linear; semicells hexagonal, angles rounded, apex truncate to retuse; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 16.5-22 µm; W: 14.8-17.3 µm; I: 3.7-4.8 µm.

Note: Present variety differs from the typical in having relatively shorter cells, about as long as broad (West & West 1908). Two different phenotypic expressions of *Cosmarium rectangulare* var. *hexagonum* were detected during the present study. Some individual specimens had smaller length:width ratio and retuse apex, similar to the type illustrated in West & West (1908), and a single specimen was longer than the others and had a very wide and truncate apex.

Present material was gathered both from lentic and lotic environments, under oligotrophic and alkaline conditions.

Material examined: BRDE25A29, BRDE25A32.

* *Cosmarium trilobulatum* Reinsch var. *abscissum* (Schmidle) Willi Krieger & Gerloff, Die Gattung *Cosmarium* 1: 99, pl. 21, fig. 1. 1962.

Figure 33

Cell 1.2-1.3 times longer than broad, median constriction deep, sinus linear; semicell trapezoidal, apex truncate, basal angles rounded; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 17.8-28.1 µm; W: 13.2-21.1 µm; I: 7.6-5 µm.

Note: According to Bicudo *et al.* (2019), *Cosmarium trilobulatum* var. *abscissum* differs from the typical variety in its broader lateral lobes and a wide apical lobe, matching with the present specimens examined. This variety is similar to *Cosmarium pseudoretusum* F.Ducellier, but differs by having less prominent apex.

Present material was collected exclusively from lentic environments, living under ultraoligotrophic-supertrophic and alkaline conditions.

Material examined: BRDE18A03, BRDE18A04, BRDE19A04, BRDE25A08, BRDE18A14, BRDE18A15, BRDE25A29, BRDE19A16.

** *Cosmarium trilobulatum* Reinsch var. *tumidum* Willi Krieger & Gerloff, Gattung *Cosmarium* 1: 101, pl. 21, fig. 8. 1962.

Figure 34

Cell 1.2-1.3 times longer than broad, median constriction deep, sinus linear; semicells trapezoidal, apex truncate, thickened, angles rounded to slightly truncate; cell wall smooth, chloroplast 1, pyrenoid 1 per semicell. L: 26.3-28.7 μm ; W: 20.3-22.9 μm ; I: 6.7-7.3 μm .

Note: This variety differs from the typical of the species in its greater dimensions, the inflated region observed in the apical view, and the slight truncate angles (Krieger & Gerloff 1962). Material presently studied was slightly smaller than that original in Krieger & Gerloff (1962).

Present material occurred in lotic and lentic environments, under oligotrophic to supertrophic and circumneutral to alkaline conditions. This is the first record of the occurrence of *Cosmarium trilobulatum* var. *tumidum* in Brazil.

Material examined: BRDE21A01, BRDE25A06, BRDE00A04, BRDE00A05, BRDE21A05, BRDE22A04, BRDE00A06, BRDE21A06, BRDE22A05, BRDE26A06, BRDE00A07, BRDE21A07, BRDE22A06, BRDE26A07, BRDE00A08, BRDE22A07, BRDE00A09, BRDE21A09, BRDE17A10, BRDE00A10, BRDE22A09, BRDE26A10, BRDE00A11, BRDE22A10, BRDE26A11, BRDE25A24, BRDE00A13, BRDE21A13, BRDE22A12, BRDE26A13, BRDE21A14, BRDE22A13, BRDE22A14.

Final remarks

Cosmarium abbreviatum var. *minus* was the only species occurring in all kinds of environments studied, present in lakes, lagoons and rivers. Most species occurred in lentic environments, only seven in lotic systems, and *Cosmarium impressulum* was the only species that showed preference for the latter kind of habitat. The most frequent species was *Cosmarium granatum* var. *granatum*, followed by *C. abbreviatum* var. *minus*. Among the environments studied, Limão lake and Monsarás lagoon presented the greatest diversity, the first with 14 and the last one with 12 out of the 22 species identified during the present inventory. Concerning pH preference, differing from the idea that genus *Cosmarium* inhabit most preferably clean and acid waters (Bicudo & Menezes 2017), only four taxa occurred in acidic conditions, most species living in circumneutral and alkaline conditions. Most species exhibited a wide distribution in terms of trophic state, with 10 species occurring up to mesotrophic conditions and 12 above eutrophic conditions. Values of total iron varied from 0.1 to 7.1 mg L^{-1} in studied systems, the values for each sampling station are detailed in Zorzal-Almeida *et al.* (2021). The great majority of the species occurred in conditions bellow 4.5 mg L^{-1} , and almost half of them bellow 1.5 mg L^{-1} . Two species occurred under conditions of high levels of total iron when compared to the others, which were *Cosmarium norimbergense* var. *depressum* (5.7 mg L^{-1}) and *Cosmarium granatum* var. *granatum* (7.1 mg L^{-1}). Finally, current study contributed with

ecological and taxonomic information to genus *Cosmarium* Corda, especially for tropical freshwater environments, and reinforces the need for more floristic studies in Brazil with ecological information, mainly in some neglected States such as Espírito Santo State.

Acknowledgments

Present study is part of the Aquatic Biodiversity Monitoring Program, Ambiental Area I, established by the Technical-Scientific Agreement, DOU number 30/2018, between FEST and Renova Foundation. We are grateful to Prof. Gilberto F. Barroso from LimnoLab/UFES, for providing data from pH, phosphorus and chlorophyll-*a* and the map development.

Author contributions

Bruna Fadul-Souza: Substantial contribution in the concept and design of the study; Contribution to data collection; Contribution to data analysis and interpretation; Contribution to manuscript preparation.

Carlos Eduardo de Mattos Bicudo: Contribution to data analysis and interpretation; Contribution to critical revision, adding intellectual content.

Geraldo José Peixoto Ramos: Contribution to data analysis and interpretation; Contribution to critical revision, adding intellectual content.

Karoline Barros de Souza: Contribution to data collection; Contribution to critical revision.

Mateus Sardi Trancoso: Contribution to data collection; Contribution to critical revision.

Stéfano Zorzal-Almeida: Contribution in the concept and design of the study; Contribution to manuscript preparation; Contribution to critical revision, adding intellectual content.

Valéria de Oliveira Fernandes: Contribution to critical revision.

Conflicts of interest

There is no conflict of interest.

Literature cited

- Aquino, C.A.N., Bueno, N.C. & Menezes, V.C.D.** 2014. Desmidióflórula (Zygnemaphyceae, Desmidiaceae) do rio Cascavel, oeste do Estado do Paraná, Brasil. *Hoehnea* 41(3): 365-392.
- Aquino, C.A.N., Bueno, N.C., Servat, L.C. & Bortolini, J.C.** 2016. New records of *Cosmarium* Corda ex Ralfs in lotic environment, adjacent to the Iguaçu National Park, Paraná State, Brazil. *Hoehnea* 43(4): 669-688.
- Bansod, V.I. & Patil, N.H.** 2019. Diversity of genus *Cosmarium* from Bodalkasa dam, Gondia district, Maharashtra, India. *Journal of Emerging Technologies and Innovative Research* 6: 124-130.
- Barroso, G.F., Garcia, F.D.C., Gonçalves, M.A., Martins, F.D.O., Venturini, J.C., Sabadini, S. & Bertoldi, L.** 2012. Estudos integrados no sistema lacustre do baixo rio Doce (Espírito Santo). *Anais do 1º Seminário Nacional de Gestão Sustentável de Ecossistemas Aquáticos: Complexidade, Interatividade e Ecodesenvolvimento, Arraial do Cabo*, pp. 126-132.
- Bicudo, C.E.M.** 1988. Polymorphism in the Desmid *Cosmarium abbreviatum* var. *minus* (Zygnemaphyceae) and its taxonomic implications. *Acta Botanica Brasilica* 2(1-2): 1-6.
- Bicudo, C.E.M., Araújo, A. & Biolo, S.** 2019. Flora Ficológica do Estado de São Paulo, 4(3): Zygnemaphyceae. Editora RiMa, São Carlos.
- Bicudo, C.E.M. & Menezes, M.** 2017. Gêneros de algas de águas continentais do Brasil. Editora RiMa, São Carlos. 3ª edição.
- Biolo, S. & Bicudo, C.E.M.** 2018. Checklist of the genus *Cosmarium* (Zygnematophyceae, Streptophyta) from Brazil. *Biodiversity International Journal* 2(5): 452-454.
- Biolo, S., Bueno, N.C., Siqueira, N.S. & Moresco, C.** 2013. New records of *Cosmarium* Corda ex Ralfs (Desmidiaceae, Zygnematophyceae) in a tributary of the Itaipu Reservoir, Paraná, Brazil. *Acta Botanica Brasilica* 27(1): 1-12.
- Biswas, S. & Nweze, N.O.** 1990. Phytoplankton of Ogelube Lake, Opi, Anambra State, Nigeria. *Hydrobiologia* 199(1): 81-86.
- Borge, O.** 1903. Die Algen der ersten Regnellschel Expedition 2: Desmidiaceen. *Arkiv für Botanik* 1: 71-138.
- Bortolini, J.C., Moresco, C., Biolo, S., Siqueira, N.S. & Bueno, N.C.** 2010. *Cosmarium* Corda ex Ralfs (Desmidiaceae) em um lago artificial urbano, Paraná, Brasil. *Revista Brasileira de Biociências* 8(3): 229-237.
- Brébisson, L.A.** 1856. Liste des desmidiées obserbées en Basse-Normandie. JB Baillièere. Paris.

- Burd, B.J.** 2002. Evaluation of mine tailing's effects on a benthic marine infaunal community over 29 years. *Marine Environmental Research* 53(5): 481-519.
- Burle, E.C., Gondra, R.M., Oliveira, I.B. & Oliveira, M.I.U.** 2020. Uma lagoa dunar e os primeiros registros de Desmidiaceae (Zygnematophyceae) em Sergipe, Brasil. *Journal of Neotropical Biology* 17(2): 162-171.
- Carmo, F.F., Kamino, L.H.Y., Junior, R.T., de Campos, I.C., Carmo, F.F., Silvino, G. & Pinto, C.E.F.** 2017. Fundão tailings dam failures: the environment tragedy of the largest technological disaster of Brazilian mining in global context. *Perspectives in ecology and conservation* 15(3): 145-151.
- Carvalho, J.K., Moresco, C., & Burack, C.** 2015. *Cosmarium* (Zygnemaphyceae) de um lago do Parque Municipal Joaquim Teodoro de Oliveira, Campo Mourão-PR. *SaBios-Revista de Saúde e Biologia* 10(2): 1-13.
- Coesel, P.F. & Meesters, K.J.** 2007. Desmids of the lowlands: Mesotaeniaceae and Desmidiaceae of the European lowlands. KNNV publishing, Zeist.
- Correa, J.A., Castilla, J.C., Ramírez, M., Varas, M., Lagos, N., Vergara, S. & Brown, M.T.** 1999. Copper, copper mine tailings and their effect on marine algae in northern Chile. In: *Developments in Hydrobiology* (vol. 137). Sixteenth International Seaweed Symposium, Cebu City, pp. 571-581.
- Croasdale, H.** 1956. Freshwater algae of Alaska, 1: some desmids from the interior, 2: *Actinotaenium*, *Micrasterias* and *Cosmarium*. *Transactions of the American Microscopical Society* 75(1): 1-70.
- Delazari-Barroso, A., Sant'Anna, C.L. & Senna, P.A.C.** 2007. Phytoplankton from Duas Bocas Reservoir, Espírito Santo State, Brazil (except diatoms). *Hoehnea* 34(2): 211-229.
- Dhande, J.S. & Jawale, A.K.** 2009. Genus *Cosmarium* Corda from Hartala Lake district, Jalgaon, Maharashtra. *Shodh, Samiksha Aur Mulyankan* 2: 196-198.
- Duque, S.R. & Donato R., J.C.** 1996. Primeros registros de *Actinotaenium* y *Cosmarium* (Desmidiaceae) en lagos de la orilla colombiana del río Amazonas. *Caldasia* 18(2): 203-210.
- Elster, J., Syobody, J., Komárek, J. & Marvan, P.** 1997. Algal and cyanoprocaryote communities in a glacial stream, Sverdrup Pass, 79 N, Central Ellesmere Island, Canada. *Archiv fur Hydrobiologie, supplement* 119: 57-93.
- Estrela, L.M.B., Fonseca, B.M. & Bicudo, C.E.M.** 2011. Desmídias perifíticas de cinco lagoas do Distrito Federal, Brasil, 1: gênero *Cosmarium* Corda ex Ralfs. *Hoehnea* 38(4): 527-552.
- Felisberto, S.A. & Rodrigues, L.** 2004. Periphytic desmids in Corumbá reservoir, Goiás, Brazil: genus *Cosmarium* Corda. *Brazilian Journal of Biology* 64(1): 141-150.

- Felisberto, S.A. & Rodrigues, L.** 2010. *Cosmarium* (Desmidiaceae, Zygnemaphyceae) da fitoflora perifítica do reservatório de Rosana, bacia do rio Paranapanema, Paraná/São Paulo, Brasil. *Hoehnea* 37(2): 267-292.
- Hatje, V., Pedreira, R.M., de Rezende, C.E., Schettini, C.A.F., Souza, G.C., Marin, D.C. & Hackspacher, P.C.** 2017. The environmental impacts of one of the largest tailing dam failures worldwide. *Scientific reports* 7(1): 1-13.
- Islam, A.N. & Irfanullah, H.M.** 2006. Hydrobiological studies within the tea gardens at Srimangal, Bangladesh, 5: desmids (*Euastrum*, *Micrasterias*, *Actinotaenium* and *Cosmarium*). *Bangladesh Journal of Plant Taxonomy* 13(1): 1-20.
- Kanetsuna, Y.** 2004. New and interesting desmids (Zygnematales, Chlorophyceae) collected from Brazil and Argentina. *Phycological Research* 52: 160-167.
- Krieger, W.** 1932. Die Desmidiaceen der deutschen limnologischen sunda-expedition. *Archiv für Hydrobiologie, Suppl.* 11: 129-230.
- Krieger, W. & Gerloff, J.** 1962. Die Gattung *Cosmarium*, Lieferung 1. Verlag von J. Cramer, Weinheim.
- Lavoie, I., Lavoie, M. & Fortin, C.** 2012. A mine of information: benthic algal communities as biomonitors of metal contamination from abandoned tailings. *Science of the Total Environment* 425: 231-241.
- Lee, O.M.** 2015. Additions to the six taxa of the genus *Cosmarium* (Desmidiaceae, Charophyta) in Korea. *Journal of Ecology & Environment* 38(4): 629-636.
- Lundell, P.M.** 1871. De Desmidiaceis, quae in Suecia inventae sunt, observationes criticae. *Nova Acta Regiae Societatis Scientiarum Upsaliensis, ser. 3*, 8(2): 1-100.
- Menezes, V.C.D., Bueno, N.C., Bortolini, J.C., Biolo, S. & Siqueira, N.S.** 2011. O gênero *Cosmarium* Corda ex Ralfs (Desmidiaceae) no Reservatório de Itaipu, PR, Brasil. *Hoehnea* 38(3): 483-493.
- Messikommer, E.** 1942. Beitrag zur Kenntnis der Algenflora und Algenvegetation des Hochgebirges um Davos. *Beiträge zur geobotanischen Landesaufnahme der Schweiz* 24: 1-452.
- Moresco, G.A., Paula, A.C.M., Bortolini, J.C., Jati, S., Reis, L.M. & Rodrigues, L.C.** 2015. Zygnemaphyceae em um lago de várzea na planície de inundação do alto rio Paraná: gêneros *Closterium*, *Cosmarium*, *Euastrum*, *Micrasterias* e *Pleurotaenium*. *Iheringia Série Botânica* 70(1): 143-155.
- Nordstedt, C.F.O. & Wittrock, V.** 1876. Desmidieae et Oedogonieae ab O. Nordstedt in Italia et Tyrolia collectae, quas determinaverunt. *Vetenskaps-akademiens Förhandlingar* 33(6): 25-56.
- Oliveira, I.B.D., Bicudo, C.E.M. & Moura, C.W.N.** 2010. Contribuição ao conhecimento de *Cosmarium* Corda ex Ralfs (Desmidiaceae, Zygnematophyceae) para a Bahia e o Brasil. *Hoehnea* 37(3): 571-600.
- Oliveira, I.B.D., Bicudo, C.E.M. & Moura, C.W.N.** 2011. New records of *Cosmarium* (Desmidiaceae) to Brazil. *Phytotaxa* 26: 25-38.

- Oliveira, I.B.D., Bicudo, C.E.M. & Moura, C.W.N.** 2016. Adição de registros de *Cosmarium* Corda ex Ralfs (Conjugatophyceae, Desmidiaceae) para a região Nordeste do Brasil. *Hoehnea* 43(2): 217-236.
- Osório, N.C., Polinario, A.M., Dunck, B., Adame, K.L., Carapunarla, L., Junqueira, M.G. & Rodrigues, L.** 2018. Periphytic *Cosmarium* (Zygnematophyceae, Desmidiaceae) in lentic environments of the Upper Paraná River floodplain: taxonomy and ecological aspects. *Acta Limnologica Brasiliensia* 30: e203.
- Ralfs, J.** 1848. *The British Desmidiaceae*. Reeve, Benham and Reeve. London.
- Ramos, G.J.P., Bicudo, C.E.M. & Moura, C.W.N.** 2018. Some new, rare and interesting desmids from bromeliad phytotelmata in Brazil. *Phytotaxa* 346(1): 59-77.
- Ramos, G.J.P., Santos, M.A.D., & Moura, C.W.N.** 2021. How hidden is the diversity of the genus *Cosmarium* (Desmidiaceae) in the Brazilian Caatinga? *Acta Botanica Brasilica*, 35: 188-214.
- Rodrigues, G.A.G. & Moresco, C.** 2021. *Cosmarium* (Zygnemaphyceae) from rural artificial lakes of Campo Mourão, PR. *Revista de Saúde e Biologia* 16: 1-22.
- Salonen, V.P., Tuovinen, N. & Valpola, S.** 2006. History of mine drainage impact on Lake Orijärvi algal communities, SW Finland. *Journal of Paleolimnology* 35(2): 289-303.
- Saturnino, M., Sophia, M.G. & Melo, S.** 2020. O gênero *Cosmarium* Corda ex Ralfs (Zygnemaphyceae, Desmidiaceae) de um lago de inundação da bacia do rio Tapajós, Amazonia, Brasil. *Amazonian Biota* 10(3): 58-64.
- Scott, A.M. & Grönblad, R.** 1957. New and interesting desmids from the southeastern United States. *Acta Societatis Scientiarum Fennicae* 2: 1-62.
- Silva, S.R.V.F. & Cecy, I.I.T.** 2004. Desmídias (Zygnemaphyceae) da área de abrangência da Usina Hidrelétrica de Salto Caxias, Paraná, Brasil, 1: gênero *Cosmarium*. *Iheringia: sér. Bot.* 59(1): 13-26.
- Šťastný, J.** 2009. The desmids of the Swamp Nature Reserve (North Bohemia, Czech Republic) and a small neighbouring bog: species composition and ecological condition of both sites. *Fottea* 9(1): 135-148.
- Šťastný, J.** 2010). Desmids (Conjugatophyceae, Viridiplantae) from the Czech Republic; new and rare taxa, distribution, ecology. *Fottea* 10(1): 1-74.
- Taniguchi, G.M., Peres, A.C., Senna, P.A.C. & Compère, P.** 2003. The desmid genera *Cosmarium*, *Actinotaenium* and *Cosmocladium* from an oxbow lake, Jataí Ecological Station (Southeastern Brazil). *Systematics and Geography of Plants* 73(1): 133-159.
- West, W. & West, G.S.** 1895. A contribution to our knowledge of the freshwater Algæ of Madagascar. *Transactions of the Linnean Society of London: sér. 2, Botany* 5(2): 41-90.
- West, W. & West, G.S.** 1902. A contribution to the freshwater Algae of Ceylon. *Transactions of the Linnean Society of London: sér. 2, Botany* 6(3): 123-215.
- West, W. & West, G.S.** 1908. A monograph of the British Desmidiaceae III. The Ray Society, London.

Special Article of Flora perifítica
<https://doi.org/10.1590/2236-8906-22/2022>

Zorzal-Almeida, S., Souza, K.B., Santana, L.M., Endlich, B.G., Fadul-Souza, B., Jesus, D.B., Balinhas, I.M., Trancoso, M.S., Scardua-Filho, M.A. & Fernandes, V.O. 2021. Periphytic algal flora of the lower Doce river basin after ore tailings flow, Espírito Santo State, Brazil: Sampling design and methods. *Hoehnea*, 48: e182021.

Editora Associada: Giselle Marquardt
Submissão: 31/05/2022
Aceito: 02/08/2022

Table 1. Distribution, frequency occurrence and environmental conditions of the non-ornamented *Cosmarium* Corda taxa in low Doce river, Espírito Santo State, Brazil. RF: relative frequency. C: categorization - LF: little frequent, R: rare. Fe: total iron. Trophic State - ult: ultraoligotrophic, oli: oligotrophic, mes: mesotrophic, eu: eutrophic, sup: supereutrophic, hyp: hypereutrophic, ac: acid, neu: circumneutral, alk: alkaline.

Taxon	River			Lagoon			Lake		Frequency		Environmental condition		
	Doce	Guandu	Areal	Areão	Monsarás	Limão	Nova	Juparanã	RF (%)	C	Trophic State	pH	Fe (mg L ⁻¹)
<i>Cosmarium abbreviatum</i> Raciborski var. <i>minus</i> (West & G.S.West) Willi Krieger & Gerloff	x	x	x	x	x	x	x	x	21.4	LF	oli-hyp	ac-alk	0.2-3.7
<i>Cosmarium angulosum</i> Brébisson	---	---	---	---	x	---	---	---	0.4	R	oli	alk	2.7
<i>Cosmarium baileyi</i> Wolle	---	---	---	---	x	---	x	---	1.1	R	mes-hyp	neu-alk	0.2-0.7
<i>Cosmarium clepsydra</i> Nordstedt var. <i>dissimile</i> (Raciborski) Willi Krieger & Gerloff	---	---	---	---	---	x	---	---	0.4	R	oli-eu	alk	0.7
<i>Cosmarium contractum</i> Kirchner var. <i>minutum</i> (Delponte) Coesel	---	---	---	---	---	x	x	x	8.4	R	ult-eu	ac-alk	0.1-1.3
<i>Cosmarium contractum</i> Kirchner var. <i>sparsipunctatum</i> Kurt Förster ex Kurt Förster	---	---	---	---	---	x	x	---	0.8	R	ult-eu	neu	0.4-0.6
<i>Cosmarium difficile</i> Lütkemüller	---	---	x	---	x	x	---	---	4.6	R	oli-eu	neu-alk	0.2-3.0
<i>Cosmarium galeritum</i> Nordstedt var. <i>subtumidum</i> Borge	---	---	---	---	x	x	x	x	3.8	R	ult-eu	neu	0.1-2.7
<i>Cosmarium granatum</i> Brébisson ex Ralfs var. <i>granatum</i>	x	x	---	---	x	x	---	---	29.0	LF	ult-sup	ac-alk	0.2-7.1
<i>Cosmarium impressulum</i> Elfving	x	---	---	---	---	---	---	---	0.4	R	mes	alk	1.4
<i>Cosmarium laeve</i> Rabenhorst var. <i>laeve</i>	x	x	x	---	---	x	---	x	20.6	LF	ult-sup	neu-alk	0.2-4.3
<i>Cosmarium lundellii</i> Delponte var. <i>madagascariense</i> West & G.S. West	---	---	---	---	---	x	---	---	0.4	R	mes	alk	0.4
<i>Cosmarium moniliforme</i> (Turpin) Ralfs	---	---	---	---	x	---	---	---	0.8	R	oli-mes	alk	2.4-3.4
<i>Cosmarium norimbergense</i> Reinsch var. <i>depressum</i> (West & G.S.West) Willi Krieger & Gerloff	---	---	---	---	x	---	---	---	1.1	R	ult-oli	neu-alk	0.7-5.7
<i>Cosmarium ocellatum</i> Eichler & Gutwinski	---	---	---	---	---	x	---	---	1.5	R	mes	neu	0.9
<i>Cosmarium pachydermum</i> P.Lundell var. <i>aethiopicum</i> (West & G.S.West) West & G.S.West	x	---	---	---	---	x	x	---	0.4	R	ult-mes	alk	0.2-1.4
<i>Cosmarium polygonum</i> (Nägeli) W.Archer var. <i>depressum</i> Messikommer	---	---	---	x	---	---	---	---	1.5	R	mes	neu-alk	1.9-2.0
<i>Cosmarium pseudoconnatum</i> Nordstedt	x	---	---	---	x	x	x	x	10.3	LF	oli-sup	ac-alk	0.1-3.8
<i>Cosmarium pseudopyramidatum</i> P.Lundell	---	---	---	---	---	x	---	---	0.4	R	oli	alk	0.7

Taxon	River			Lagoon			Lake			Frequency		Environmental condition		
	Doce	Guandu	Areal	Areão	Monsarás	Limão	Nova	Juparanã	RF (%)	C	Trophic State	pH	Fe (mg L ⁻¹)	
<i>Cosmarium rectangulare</i> Grunow var. <i>hexagonum</i> (Delponte) West & G.S. West	---	---	---	---	x	---	---	---	0.8	R	oli	alk	2.7-3.4	
<i>Cosmarium trilobulatum</i> Reinsch var. <i>abscissum</i> (Schmidle) Willi Krieger & Gerloff	---	---	---	---	x	x	x	---	3.1	R	ult-sup	alk	0.2-2.7	
<i>Cosmarium trilobulatum</i> Reinsch var. <i>tumidum</i> Willi Krieger & Gerloff	x	x	---	---	x	---	---	---	16.8	LF	oli-sup	neu-alk	0.8-3.7	




Figure 1. *Cosmarium abbreviatum* Raciborski var. *minus* (West & G.S.West) Willi Krieger & Gerloff. 2. *Cosmarium angulosum* Brébisson. 3. *Cosmarium baileyi* Wolle. 4-6. *Cosmarium clepsydra* Nordstedt var. *dissimile* (Raciborski) Willi Krieger & Gerloff. 7, 8. *Cosmarium contractum* Kirchner var. *minutum* (Delponte) Coesel. 9, 10. *Cosmarium contractum* Kirchner var. *sparsipunctatum* Kurt Förster ex Kurt Förster. 11, 12. *Cosmarium difficile* Lütkenmüller. 13, 14. *Cosmarium galeritum* Nordstedt var. *subtumidum* Borge. 15, 16. *Cosmarium granatum* Brébisson ex Ralfs var. *granatum*. 17. *Cosmarium impressulum* Elfving. 18, 19. *Cosmarium laeve* Rabenhorst var. *laeve*. 20, 21. *Cosmarium lundellii* Delponte var. *madagascariense* West & G.S. West. 22. *Cosmarium moniliforme* (Turpin) Ralfs. 23. *Cosmarium norimbergense* Reinsch var. *depressum* (West & G.S.West) Willi Krieger & Gerloff. 24. *Cosmarium ocellatum* Eichler & Gutwinski. 25. *Cosmarium pachydermum* P.Lundell var. *aethiopicum* (West & G.S.West) West & G.S.West. 26, 27. *Cosmarium pseudoconnatum* Nordstedt. 28, 29. *Cosmarium polygonum* (Nägeli) W.Archer var. *depressum* Messikommer. 30. *Cosmarium pseudopyramidatum* P.Lundell. 31, 32. *Cosmarium rectangulare* Grunow var. *hexagonum* (Delponte) West & G.S.West. 33. *Cosmarium trilobulatum* Reinsch var. *abscissum* (Schmidle) Willi Krieger & Gerloff. 34. *Cosmarium trilobulatum* Reinsch var. *tumidum* Willi Krieger & Gerloff. Scale bar = 10 µm.

CARTA DE AUTORIZAÇÃO DE PUBLICAÇÃO NO PORTAL DE PREPRINTS DO SCIELO

Ao Comitê Editorial de HOEHNEA

Declaro, em meu próprio nome e nos dos demais Autores, que concordo com a publicação do Artigo Aceito pelo Corpo Editorial de Hoehnea, intitulado “Periphytic algal flora of the Low Doce River, Espírito Santo State, Brazil, after ore tailings flow, 1: smooth cell walled *Cosmarium* (Zygnematophyceae, Desmidiaceae)” de autoria de Bruna Fadul-Souza, Karoline Barros de Souza, Mateus Sardi Trancoso, Stéfano Zorzal-Almeida, Carlos Eduardo de Mattos Bicudo, Geraldo José Peixoto Ramos & Valéria de Oliveira Fernandes no Portal de Preprints do SciELO Brasil (<https://preprints.scielo.org/index.php/scielo/preprints/section/biological>).

Declaro, ainda, que o referido artigo é original, sendo que o conteúdo não foi ou não está sendo considerado para publicação em outro periódico, quer seja no formato impresso e/ou eletrônico.

Campos dos Goytacazes, 12 de agosto de 2022.



BRUNA FADUL DE SOUZA

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.