

Publication status: Preprint has been published in a journal as an article
DOI of the published article: <https://doi.org/10.1590/2236-8906e392022>

Occurrence and distribution of the heterotrophic straminipiles from Brazilian Atlantic Rainforest areas

Carmen Lidia Amorim Pires-Zottarelli; Ana Lucia de Jesus, Débora Rodrigues da Silva Colombo,
Poliana de Oliveira Ventura, Marcela Castilho Boro

<https://doi.org/10.1590/2236-8906-39/2022>

Submitted on: 2023-03-01

Posted on: 2023-03-01 (version 1)

(YYYY-MM-DD)

**Occurrence and distribution of the heterotrophic straminipiles from Brazilian Atlantic
Rainforest areas**

Carmen Lidia Amorim Pires-Zottarelli^{1,2}, Ana Lucia de Jesus¹, Débora Rodrigues da Silva Colombo¹,
Poliana de Oliveira Ventura¹ and Marcela Castilho Boro¹

Título resumido: Heterotrophic straminipiles from Brazilian Atlantic Rainforest

Carmen Lidia Amorim Pires-Zottarelli: <https://orcid.org/0000-0003-1710-057X>

Ana Lucia de Jesus: <https://orcid.org/0000-0002-7466-7030>

Débora Rodrigues da Silva Colombo: <https://orcid.org/0000-0003-1263-9509>

Poliana de Oliveira Ventura: <https://orcid.org/0000-0001-5686-7728>

Marcela Castilho Boro: [https:// orcid.org/0000-0002-0555-8957](https://orcid.org/0000-0002-0555-8957)

¹ Instituto de Pesquisas Ambientais - Unidade Jardim Botânico, Secretaria de Meio Ambiente, Infraestrutura e Logística, Avenida Miguel Estéfno, 3687, Água Funda, 04301-902 São Paulo, SP, Brazil

² Corresponding author: zottarelli@uol.com.br

ABSTRACT – (Occurrence and distribution of the heterotrophic straminipiles from Brazilian Atlantic Rainforest areas). We present herein the occurrence and distribution of the heterotrophic straminipiles collected from soil and water, submerged leaves and surface sediment from freshwater bodies at “Mosaico de Unidades de Conservação Juréia-Itatins”, a preserved conservation unit of the Brazilian Atlantic Rainforest. From 180 collected samples, we isolated 41 taxa belonging to Hyphochytriomycota and Oomycota using the multiple baiting technique, with several new taxa to the science, new occurrences to São Paulo State or Brazil, and all new citation to the studied area. These taxa were identified based on morphological and/or molecular data, with several genic regions made available in GenBank. The calculated community structure of these zoosporic organisms showed a diverse community, with few dominant taxa, without significant differences ($P \leq 0.05$) among the different types of samples.

Keywords: diversity, ecology, Hyphochytriomycota, Oomycota, taxonomy

RESUMO – (Ocorrência e distribuição dos estramenópilos heterotróficos de áreas de Mata Atlântica Brasileira). Apresentamos aqui a ocorrência e a distribuição dos estramenópilos heterotróficos coletados de solo, e de água, folhas submersas e sedimento superficial de corpos de água doce no “Mosaico de Unidades de Conservação Juréia-Itatins”, uma unidade de conservação preservada de Mata Atlântica brasileira. Das 180 amostras coletadas, isolamos 41 táxons pertencentes aos filos Hyphochytriomycota e Oomycota, usando a técnica de iscagem múltipla, com vários deles novos táxons para a ciência, novas ocorrências para o estado de São Paulo ou Brasil, e todos primeira ocorrência para o local estudado. Os táxons foram identificados com base em dados morfológicos e/ou moleculares, com várias regiões gênicas disponibilizadas no GenBank. O cálculo da estrutura da comunidade mostrou uma comunidade diversa, com poucos táxons dominantes, sem diferença significativa ($P \leq 0.05$) entre os diferentes tipos de amostras. **Palavras-chave:** diversidade, ecologia, Hyphochytriomycota, Oomycota, taxonomia

Introduction

The heterotrophic straminipiles are fungal-like organisms inserted within the Kingdom Straminipila, in the supergroup SAR, together with the Alveolata and Rhizaria (Adl *et al.* 2019). It is represented by the phyla Hyphochytriomycota, Labyrinthulomycota and Oomycota that are characterized by their heterokont, predominantly biflagellate zoospores. Members of these phyla are widespread and can be found specially as saprotrophs of different kinds of substrates, in freshwater, marine and terrestrial

ecosystems. Some of them are important parasites of a variety of host organisms, such as algae, animals, fungi, oomycetes and plants (Beakes *et al.* 2014, Beakes & Thines 2017, Marano *et al.* 2017).

These phyla have similarities with the true zoosporic fungi, like as osmotrophic nutrition, flagellate spores (zoospores) and ecological niches, however, molecular studies have shown that the heterotrophic straminipiles are phylogenetically distant from these fungi (Beakes *et al.* 2012, Beakes & Thines 2017, Marano *et al.* 2017).

Hyphochytriomycota and Oomycota were studied in the “Mosaico de Unidades de Conservação Juréia-Itatins” from 2016 to 2017, an important Atlantic Rainforest area located in São Paulo State, Brazil, with the aim to assess the species composition and community structure of these phyla from soil and freshwater samples. Although several studies have already been carried out in Brazilian Atlantic Rainforest (Beneke & Rogers 1962, 1970, Rogers *et al.* 1970, Lyra & Milanez 1974, Pires-Zottarelli *et al.* 1995, 1996, Schoenlein-Crusius & Milanez 1998, Schoenlein-Crusius *et al.* 2006, Pires-Zottarelli & Rocha 2007, Miranda & Pires-Zottarelli 2008, 2012, Jesus *et al.* 2013, 2015, 2016a, b, Rocha *et al.* 2016a, b, 2018, and others), there was no study with these phyla in the studied area.

Material and Methods

Studied area, sampling and laboratory analyses - From August/2016 to October/2017, samples of soil, as well as water, submerged leaves and surface sediments from freshwater bodies were collected at “Mosaico de Unidades de Conservação Juréia-Itatins”, São Paulo State, Brazil (Figure 1). The aliquots of these samples were plated and baited using cellulosic, keratinous and chitinous substrates according to the technique described in Milanez (1989). These samples were incubated at room temperature (~22° C) and the baits observed after five days under a Leica DMLB2 compound microscope (Hessen, Wetzlar, Germany). The specimens observed were isolated and purified onto MP₅ culture medium (4 g maltose, 1 g peptone, 15 g agar and 1000 mL deionized water) and CMA culture medium (17 g cornmeal agar and 1000 mL deionized water) with addition of 0,05 g of streptomycin sulfate and 0,1 g of penicillin. The asexual and/or sexual structures of the obtained specimens were observed under the microscope, measured in water cultures (n=50) and identified morphologically according to specific literature, such as Scott (1961), Seymour (1970), Karling (1977), Plaats-Niterink (1981), Johnson *et al.* (2002), and original descriptions. Due to difficulties in culturing some taxa on agar media, these were purified by multiple transfers to new substrates. The identified specimens were photographed with a Leica MC170 HD camera using Leica Qwin 3.1 software (Hessen, Wetzlar, Germany), and at least one isolate was

incorporated at CCIBt (“Coleção de Culturas de Algas, Cianobactérias e Fungos do Instituto de Botânica” - nowadays “Instituto de Pesquisas Ambientais”) and/or Herbarium SP. Taxonomical placement of the genera followed Beakes & Thines (2017). *Aphanodictyon* and *Phragmosporangium*, genera not mentioned by these authors, followed Dick (2001) and Rocha *et al.* (2018), respectively, at family level.

DNA extraction, PCR amplification and sequencing - For DNA extraction, small fragments of culture medium with the mycelium of the specimens were transferred to erlenmeyer flasks (250 mL) containing 30 mL of MP₅ liquid medium. Cultures were incubated at ~22 C for 5-10 days, after which the biomass was transferred to 2 mL microfuge tubes. The tubes were centrifuged for 15 min at 13000 rpm, the supernatant removed, and the resulting pellet was used for DNA extraction. The pellets were treated according to the protocol described in the PureLink Genomic DNA kit (Invitrogen). Partial LSU, SSU, and ITS1-5.8S-ITS2 regions of the nuclear rDNA locus were amplified using the LR0R/LR6-O (Riethmüller *et al.* 2002), NS4/SR1R (White *et al.* 1990) and UN-up18S42/UN-lo28S22 (Robideau *et al.* 2011) primers, respectively, and/or COI mtDNA using OomCoxI-Levup/OomCoxI-Levlo (Robideau *et al.* 2011). Amplicons were purified using the protocol in Schmitz and Riesner (2006), and sequencing was performed using the same PCR primer sets in an ABI 3730 DNA Analyser (Life Technologies, Carlsbad, California). The sequences were edited manually using Sequencher 4.1.4. The obtained sequences were compared with similar sequences of GenBank using the BLASTn, with our sequences have been published in this database (<https://www.ncbi.nlm.nih.gov/genbank/>).

Ecological data analyses - The total frequency (TF) was calculated based on the occurrence of each taxon considering the total sampling/types of samples (16 in total - four sampling during the study period and four types of samples) according to Marano *et al.* (2008). The total abundance (TA) of the identified species was based on the number of occurrences considering the presence or absence of them in the samples. The community structure was estimated by (i) the species richness (S), (ii) Shannon diversity index $H' = -\sum_{i=1}^S p_i \log_2(p_i)$, where p_i is the abundance of the species i in the community, (iii) evenness $E = H'/H'_{\max}$, where H'_{\max} is the maximum value of the diversity for the number of present taxa, (iv) Simpson's dominance index $D = 1/\sum_{i=1}^S (p_i)^2$ based on the species abundance (Hammer *et al.* 2001, Zak & Willig 2004). The Wilcoxon test was used to verify if the abundance and diversity measured were equal between the samples (Hammer *et al.* 2001).

Results

Species composition and community structure - One hundred and eighty samples (45 water, 45 soil, 45 submerged leaves, 45 surface sediment) were analyzed. From these, 209 specimens were obtained, and 41 taxa identified belonging to the phyla Hyphochytriomycota (1) and Oomycota (40), with 38 at species level and 03 to genus level, as shown in the Table 1. Total frequency and abundance of them are also mentioned. Thirty-eight species were sequenced (ITS, SSU and LSU rDNA and/or COI mDNA) and most of the sequences available in GenBank. Twelve taxa have already been published as new species and/or new citation for Brazil or the studied area (Pires-Zottarelli *et al.* 2019, Dima *et al.* 2021, Pires-Zottarelli *et al.* 2022).

From 41 identified taxa, 21 have been occurred in water (53 occurrences), 17 in submerged leaves (40 occurrences), 18 in surface sediment (39 occurrences) and 25 in soil samples (77 occurrences). Although the species richness (S) and occurrence were higher in soil, there was no significant differences ($P \leq .05$) among the samples considering the indices (Table 2). *Achlya* sp1, *Aphanomyces stellatus*, *Phragmosporangium uniseriatum* e *Plectospira myriandra* were the species more frequent and abundant, considering all samples.

For each taxa presented below, geographic location, basionym, origin and GenBank accession numbers (not yet mentioned or published elsewhere) are provided. Images of the species not previously illustrated by our team are presented. New citations of the species for the São Paulo State and/or the studied area are mentioned herein.

KINGDOM STRAMINIPILA

PHYLUM HYPHOCHYTRIOMYCOTA

HYPHOCHYTRIALES

HYPHOCHYTRIACEAE

1. *Hyphochytrium catenoides* Karling. Am. J. Bot. 26: 512. 1939.

Figures 2 a, 2 b

Specimen examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from soil samples baited with snake skin, 27 June 2017. Culture not obtained.

Identification: This species of *Hyphochytrium* is characterized by a polycentric thallus, usually hypha-like, with a linear series of intercalary and terminal swellings and zoosporangia. The zoosporangia are normally spherical or ovoid and delimited by cross septa. The zoospores anteriorly uniflagellate with numerous lipidic granules emerge by exit tubes from the zoosporangia as a naked mass, maturing outside

the zoosporangia (Karling 1939). The characteristics of our specimen are in agreement with the original description. This species was previously mentioned in Brazil by Moreira (2006) from submerged leaves of *Tibouchina pulchra* Cogn. in Atlantic Rainforest area in São Paulo city. This is a new citation for the studied area.

PHYLUM OOMYCOTA

LEPTOLEGNIELLACEAE - *Incertae sedis*

2. *Aphanodictyon papillatum* Huneycutt ex M.W. Dick, Trans. Br. Mycol. Soc. 57: 422. 1971.

Figures 2 c, 2 d

Specimen examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from soil samples baited with snake skin, 25 Oct 2017, SP512792.

Culture not obtained.

Identification: *Aphanodictyon papillatum* grow only as saprotrophic in keratinous substrate and their pure culture has never been obtained. We tried many times in several media cultures, however no culture was obtained. It is characterized by delicate, much-branched hyphae, normally globose to subglobose zoosporangia. Their biflagellate zoospores are discharged like as *Dictyuchus* genus, leaving a net of cysts wall into the zoosporangia. The oogonia are normally globose to subglobose, some irregular, with papillate ornamentations; androgynous or diclinous antheridia; thick oospores normally 1-3 by oogonia (Huneycutt 1948). The characteristics of our specimen are in agreement with the original description. This species was previously mentioned in Brazil from soil samples in Piauí State (Rocha 2002), with their distribution in the country published by Milanez *et al.* (2007) and Rocha *et al.* (2010). This is a new citation for São Paulo State.

3. *Leptolegniella exogena* Karling. Nova Hedwigia 45: 433. 1987.

Specimen examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from soil samples baited with snake skin, 27 June 2017, SP512793, CCIBt4682. GenBank – SSU: ON860582, LSU: ON860619.

Identification: This holocarpic species presented a coarse branched occasionally constricted and septate mycelium, which is intramatrical or extramatrical in the substrate. Zoosporangia undifferentiated of vegetative mycelium. Zoospores biflagellate normally emerging in succession and forming a globular mass at the exit orifice or encysting into the sporangia. Development of the resting spores is unique for this species, where the protoplasm aggregated into the hyphae contract to form exogenous resting spores

that remain attached to the hyphae wall (Karling 1987). Our specimen presented all the important characteristics of this beautiful species. Sequences of this genus/species from were obtained for the first time from pure culture. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

4. *Leptolegniella keratinophila* Huneycutt, J. Elisha Mitchell. Sci. Soc. 68: 110. 1952.

Specimens examined: BRAZIL. SÃO PAULO: Peruíbe, Pedro de Toledo, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W; 24°21'59.1"S, 47°18'32.7"W; 24°34'21.1"S, 47°14'53.0"W, respectively, from submerged leaves of freshwater bodies and soil samples baited with snake skin, 24 Aug 2016, 27 June 2017 and 25 Oct 2017, SP512794. Culture not obtained.

Identification: This holocarpic species described by Huneycutt (1952) is characterized by a poorly developed extramatrical and an extensive intramatrical mycelium, in keratinous substrate. Zoosporangia undifferentiated of vegetative mycelium. Resting spores is typical for this species and produced within undifferentiated hyphae. Pure culture of this species has not been obtained. Our specimens are in agreement with the original description. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

5. *Leptolegniella sp1*

Specimens examined: BRAZIL. SÃO PAULO: Peruíbe and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from soil samples baited with snake skin, 24 Aug 2016 and 25 Oct 2017, CCIBt4424. Sequences are not available herein, because this isolate is being studied considering it does not fit into any previously taxa described.

PERONOSPORALES *SENSU LATO*

PYTHIACEAE *SENSU LATO*

6. *Myzocytiopsis humana* (Karling) M.W. Dick, Mycol. Res. 101: 879. 1997.

Basionym: *Lagenidium humanum* Karling

Specimens examined: BRAZIL. SÃO PAULO: Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W; 24°34'21.1"S, 47°14'53.0"W, respectively, from submerged leaves and surface sediment of freshwater bodies and soil

samples baited with snake skin, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4304. GenBank – ITS: ON860609, SSU: ON860583, LSU: ON860620.

Identification: The specimens presented septate thallus with exit tubes and numerous refractive bodies, forming disarticulated zoosporangium as mentioned in the original description of Karling (1947). Sexual reproduction was not observed. They grew in keratinous substrate (snake skin) and very well in medium culture (MP₅). It is a new citation for the studied area. It was previously mentioned for Brazil by Jesus *et al.* (2013) from water samples of the Atlantic Rainforest of São Paulo city. Illustration of the species in Jesus *et al.* (2013).

7. *Pythiogeton abundans* J.H. Huang, Chi Y Chen & Yi S. Lin, Mycoscience 54: 132. 2013.

Figure 2 e

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from water and surface sediment samples of freshwater bodies baited with *Sorghum* sp seeds, 27 Jun 2017, CCIBt4340. GenBank – ITS: ON860610.

Identification: The production of the spherical, unsymmetrical and bursiform zoosporangia, their insertion in the hyphae at right angles, the internal proliferation and the zoospores discharge of them confirmed the morphological identification of the isolate at genus level, as mentioned by Sparrow (1960). Our specimens were identified at species level taking into account the morphological and molecular data, considering that *P. abundans* not produce sexual structures as mentioned by Huang *et al.* (2013). It was previously mentioned for Brazil from freshwater samples, as well as dead trout fish in Campos do Jordão, São Paulo State (da Paixão 2021). This is a new citation for the studied area.

8. *Pythium acanthicum* Drechsler, J. Wash. Acad. Sci. 20: 408. 1930.

Figures 2 f, 2 g

Specimens examined: BRAZIL. SÃO PAULO, Peruíbe, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W; 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from surface sediment of freshwater bodies and soil samples baited with *Sorghum* sp seeds, 24 Aug 2016, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt 4309. Genbank – ITS: ON860611, LSU: ON860621.

Identification: The production of the zoosporangia, the discharge and the maturation of the zoospores into the vesicle, the oogonia with conical projections and blunt tip and monoclinous antheridia, as mentioned by Plaats-Niterink (1981), are characteristics of this species. The morphology of our

specimens, as well as the rDNA sequences, allowed us to identify the isolate as *P. acanthicum*. It is a new citation for the studied area.

9. *Pythium graminicola* Subraman, Bull. Agric. Res. Inst. Pus. 177: 1. 1928.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°33'11.1"S, 47°13'34.5"W, respectively, from surface sediment samples of freshwater bodies baited with *Sorghum* sp seeds, 14 Feb 2017 and 27 Jun 2017, CCIBt4330. GenBank - ITS: ON860612, LSU: ON860622.

Identification: This species is characterized by inflated filamentous zoosporangia, and by oogonia with predominantly monoclinal antheridia (Plaats-Niterink 1981). Our specimens were identified taking into account the morphological and molecular data. It is a new citation for the studied area. Illustration of the species in Jesus *et al.* (2013).

10. *Pythium periplocum* Drechsler, J. Wash. Acad. Sci. 20: 405. 1930.

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and submerged leaves samples of freshwater bodies baited with snake skin and *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4344, CCIBt 4420. GenBank - ITS: ON860613, LSU: ON860623.

Identification: *Pythium periplocum* is characterized by inflated filamentous zoosporangia and spines oogonia (Plaats-Niterink 1981). Our specimens produced many zoosporangia, however few oogonia. The molecular data helped us to identify the specimens. It is a new citation for the studied area. Illustration of the species in Gonçalves *et al.* (2016a).

11. *Pythium splendens* Hans Braun, J. Agric. Res. 30: 1061. 1925.

Figure 2 h

Specimen examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from surface sediment samples of freshwater bodies baited with *Sorghum* sp seeds, 25 Oct 2017, CCIBt4399. GenBank - ITS: ON860614, LSU: ON860624.

Identification: Our specimen produced globose and abundant hyphal swellings, with dark granular content, like as mentioned by Plaats-Niterink (1981). The molecular data helped us to identify the specimen. This is a new citation for São Paulo State.

PERONOSPORACEAE *SENSU LATO*

12. *Phytophthora undulata* (H.E. Petersen) M.W. Dick, Mycotaxon 35: 449. 1989.

Basionym: *Pythium undulatum* H.E. Petersen

Figures 3 a, 3 b

Specimen examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from submerged leaves samples of freshwater bodies baited with snake skin and corn leaves, 27 Jun 2017, CCIBt4303. GenBank - ITS: ON860615, LSU: ON860625.

Identification: This species is easy recognize by elongate and proliferous zoosporangia, and dark yellow chlamydospores usually intercalary in the hyphae (Plaats-Niterink 1981). Our molecular data confirm the morphological identification. It is a new citation for the studied area.

13. *Phytophthora chamaehyphon* (Sideris) Abad, de Cock, Bala, Robideau, A.M. Lodhi & Lévesque, Persoonia 34: 36. 2014.

Basionym: *Pythium chamaehyphon* Sideris

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°33'11.1"S, 47°13'34.5"W, respectively, from water, surface sediment and submerged leaves samples of freshwater bodies baited with corn leaves, onion skin and *Sorghum* sp seeds, 14 Feb 2017 and 27 Jun 2017, CCIBt4338, CCIBt4407.

Identification: Description, comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Dima *et al.* (2021).

14. *Phytophthora palingenes* (Drechsler) Abad, de Cock, Bala, Robideau & Levésque, Persoonia 34: 37. 2014.

Basionym: *Pythium palingenes* Drechsler

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from water and submerged leaves of freshwater bodies and soil samples baited with onion skin, snake skin and *Sorghum* sp seeds, 25 Oct 2017, CCIBt4397, CCIBt4428.

Identification: Comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Dima *et al.* (2021).

15. *Phytophthium vexans* (de Bary) Abad, de Cock, Bala, Robideau & Levésque, *Persoonia* 34: 37. 2014.

Basionym: *Pythium vexans* de Bary

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from soil samples baited with onion skin, snake skin and *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4383, CCIBt4398, CCIBt4422.

Identification: Comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Dima *et al.* (2021).

SAPROLEGNIALES

SAPROLEGNACEAE *SENSU LATO*

16. *Achlya catenulata* Pires-Zottar., A.L. Jesus, Marano & J.I. Souza, *Phytotaxa* 212: 223. 2015.

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and submerged leaves of freshwater bodies samples baited with shrimp exoskeleton and *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4302. GenBank - ITS: OK271001, SSU: OK235689; LSU: OK270960.

Identification: This Brazilian species was described by Jesus *et al.* (2015) from freshwater and saline water samples of the Atlantic Rainforest of São Paulo State. It is characterized by catenulate smooth-walled oogonia, usually in chains, with declinuous antheridia and eccentric oospores, which normally not mature. Our specimens are in agreement with the original description. It is a new citation for the studied area. Illustration of the species in Jesus *et al.* (2015).

17. *Achlya diffusa* J.J.V. Harv. ex T.W. Johnson, *The genus Achlya: morphology and taxonomy* 64: 1956.

Figures 3 c, 3 d

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W, from water, surface sediments and submerged leaves of freshwater bodies samples baited with *Sorghum* sp seeds, 14 Feb 2017, CCIBt4396, CCIBt4405, CCIBt4429. GenBank - ITS: OK270977, OK270979, OK270978, respectively; SSU: NA, NA, NA.

OK235673, respectively, LSU: OK270937, OK270939, OK270938, respectively; COI: OK120427, OK120429, OK120428, respectively.

Identification: This species is characterized by inflated “balloon-like” hyphal swellings, and eccentric oospores, which usually not mature (Johnson 1956). The characteristics of our specimens are in agreement with the original description. This is a new citation for São Paulo State.

18. *Achlya flagellata* Coker, Saprolegniaceae with notes on other water molds: 116. 1923.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and surface sediments of freshwater bodies and soil samples baited with shrimp exoskeleton and *Sorghum* sp seeds, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4400, CCIBt4404, CCIBt4425. GenBank - ITS: OK270982, OK270981, OK270980, respectively; SSU: NA, NA, NA, respectively; LSU: OK270942, OK270941, OK270940, respectively; COI: NA, NA, OK120430, respectively.

Identification: This species is very common in aquatic and terrestrial Brazilian ecosystems (Milanez et al. 2007). It is characterized by production of oogonia with oospheres usually not maturing, when mature are eccentric (Johnson *et al.* 2002). The molecular data confirm our morphological identification. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

19. *Achlya klebsiana* Pieters, Bot. Gaz. 60: 486. 1915.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water, surface sediments and submerged leaves of freshwater bodies samples baited with snake skin and *Sorghum* sp seeds, 14 Feb 2017 and 25 Oct 2017, CCIBt4333, CCIBt4339, CCIBt4357, CCIBt4421. GenBank - ITS: OK270984, OK270983, OK270985, OK270986, respectively; SSU: NA, OK235675, NA, NA, respectively; LSU: OK270944, OK270943, OK270945, OK270946, respectively; COI: OK120432, OK120431, OK120433, NA, respectively.

Identification: This species is characterized by oogonia with eccentric oospores and usually declinous and monoclinal antheridia, when monoclinal arising from a point remote from the oogonium to which attached as mentioned by Johnson (1956). In Johnson *et al.* (2002) this species, together with *Achlya flagellata*, are considered synonym of *Achlya debaryana*, however this is not accepted by many

specialists, including us. The molecular data obtained confirm our impression. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

20. *Achlya orion* Coker & Couch, J. Elisha Mitchell Sci. Soc. 36: 100. 1920.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water of freshwater bodies and soil samples baited with *Sorghum* sp seeds, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4310, CCIBt4384. GenBank - ITS: OK270987, OK270988, respectively; SSU: NA; LSU: OK230947, OK270948, respectively; COI: OK120434, OK120435, respectively.

Identification: This species very common in Brazilian ecosystems (Milanez *et al.* 2007) is easy recognize by usually spherical oogonia with long, slender, bent or recurved stalks, commonly androgynous or monoclinal antheridia, and eccentric oospores (Johnson *et al.* 2002). The characteristics of our specimens are in agreement with the description. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

21. *Achlya proliferoides* Coker, Saprolegniaceae with notes on other water molds 115, 1923.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°33'11.1"S, 47°13'34.5"W, respectively, from water of freshwater bodies samples baited with *Sorghum* sp seeds, 14 Feb 2017 and 27 Jun 2017, CCIBt4418, CCIBt4419. GenBank - ITS: OK270989, OK270990, respectively; SSU: NA, OK235677, respectively; LSU: OK270949, OK270950, respectively; COI: OK120436, OK120437, respectively.

Identification: This species is characterized by predominantly declinous antheridia coiling in the hyphae which may or may not bear oogonia, and eccentric oospores (Johnson *et al.* 2002). Our specimens produced all the described characteristics. It is a new citation for the studied area. Illustration of the species in Nascimento & Pires-Zottarelli (2012).

22. *Achlya truncatiformis* M.W. Dick & Mark A. Spencer, Mycol Res. 106: 557. 2002.

Figures 3 e, 3 f

Specimen examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from soil samples baited with *Sorghum* sp seeds, 25 Oct 2017, CCIBt4341. GenBank - ITS: OK270991; SSU: OK235678; LSU: OK270951; COI: OK120438.

Identification: The description of *Achlya truncatiformis* is based on *Achlya recurva sensu* Latham, which is characterized by typical truncate oogonial papillae and eccentric oospores (Spencer *et al.* 2002). Our specimen is in agreement with the description of these authors. It is a new citation for the studied area, considering that specimens of *Achlya recurva* with this type of ornamentation and eccentric oospores were mentioned for Brazilian ecosystems of Piauí and São Paulo State (Milanez *et al.* 2007).

23. *Achlya* sp 1

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water, surface sediment, submerged leaves of freshwater bodies and soil samples baited with snake skin and *Sorghum* sp seeds, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4337, CCIBt4381, CCIBt4408. Sequences are not available herein.

Identification: The specimens are being studied considering its does not fit into any previously described species.

24. *Aplanopsis terrestris* Höhnk, Veröff. Inst. Meeresf. Bremerhaven, Sonderband 1: 127. 1952.

Specimens examined: BRAZIL. SÃO PAULO: Pedro de Toledo, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W, from soil samples baited with snake skin, 14 Feb 2017, CCIBt4307. GenBank - ITS: OK270992, LSU: OK270953.

Identification: This species is characterized by no zoosporangia produced; oogonia wall usually smooth, however few papillae were observed, with centric and subcentric oospores and commonly androgynous antheridia (Johnson *et al.* 2002, Rocha *et al.* 2016a). The morphological and molecular data confirm the identification of this species. It was previously mentioned for Brazil by Rocha *et al.* (2016a) from soil samples of the Atlantic Rainforest of São Paulo State. It is a new citation for the studied area. Illustration of the species in Rocha *et al.* (2016a).

25. *Brevilegnia longicaulis* T.W. Johnson, Mycologia 42: 244. 1950.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W, from soil samples baited with *Sorghum* sp seeds, 14 Feb 2017, CCIBt4336, CCIBt4356. GenBank - ITS: OK270995, OK270994, respectively; SSU: NA, OK235684, respectively. LSU: NA, OK270955, respectively.

Identification: This species is recognized by discharge and behavior brevilegnoid, rarely dictyucooid, of the zoospores, antheridia always declinous attaching in oogonia that produce eccentric oospores. When dictyucooid behavior occurs the zoosporangium partly disintegrated and disarticulating (Johnson *et al.* 2002). It was previously mentioned for Brazil by Rocha *et al.* (2016a) from soil samples of the Atlantic Rainforest of São Paulo State. It is a new citation for the studied area. Illustration of the species in Rocha *et al.* (2016a).

26. *Brevilegnia* sp 1

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from soil samples baited with *Sorghum* sp seeds, 14 Feb 2017 and 25 Oct 2017, CCIBt4332, CCIBt4350. Sequences not available herein.

Identification: The specimens are being studied considering its does not fit into any previously described species of the genus.

27. *Dictyuchus pseudodictyon* Coker & Braxton, J. Elisha Mitchell Sci. Soc. 46: 228. 1931.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from submerged leaves of freshwater bodies samples baited with *Sorghum* sp seeds, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4377, CCIBt4378. GenBank - ITS: OK271002, OK271003, respectively; SSU: OK235690, NA; LSU: OK270961, NA, respectively.

Identification: The zoospores cysts left inside the zoosporangia, forming true and false network, declinous antheridia involving the oogonium and eccentric oospores are important features of this species (Johnson *et al.* 2002). Our morphological and molecular data of the specimens confirm the identification. It is a new citation for the studied area. Illustration of the species in Miranda & Pires-Zottarelli (2012).

28. *Leptolegnia eccentrica* Coker, J. Elisha Mitchell Sci. Soc. 42: 215. 1927.

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from soil samples baited with snake skin, 25 Oct 2017, CCIBt4403. GenBank - ITS: ON860617, LSU: ON860627.

Identification: Papillate oogonia, with papillae in variable number, monoclinal and androgynous antheridia, as well as eccentric oospores are characteristic of this species (Johnson *et al.* 2002). The

characteristics of our specimens are in agreement of the species' description. It is a new citation for the studied area. Illustration of the species in Rocha *et al.* (2018).

29. *Newbya androgyna* (W. Archer) Pires-Zottar. & S.C.O. Rocha, Mycol. Progr. 17: 696. 2018.

Basionym: *Saprolegnia androgyna* W. Archer

Specimens examined: BRAZIL. SÃO PAULO, Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water of freshwater bodies and soil samples baited with *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4306. GenBank - ITS: OK271005; SSU: OK235692; LSU: OK270963.

Identification: Rare terminal or intercalary zoosporangia, filiform and/or naviculate oogonia with papillate ornamentation, subcentric oospores, and androgynous antheridia are important features of this species (Johnson *et al.* 2002, Rocha *et al.* 2018). The characteristics of our specimens are in agreement with the descriptions of these authors. It is a new citation for the studied area. Illustration of the species in Rocha *et al.* (2018).

30. *Pythiopsis irregularis* R.L. Szym., Mycotaxon 92: 2. 2005.

Figures 3 g, 3 h

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°21'59.1"S, 47°18'32.7"W and 24°33'11.1"S, 47°13'34.5"W, respectively, from soil samples baited with snake skin, 14 Feb 2017 and 27 Jun 2017, CCIBt4308, CCIBt4311. GenBank - ITS: OK271009; SSU: OK235700; LSU: OK270967.

Identification: This species produces oogonia with short, broad, conspicuous papillae or bullations, centric and subcentric oospores and androgynous antheridia (Johnson *et al.* 2002). The characteristics of our specimens are in agreement of the original description, with molecular data corroborating the morphological identification. It is a new citation for the studied area.

31. *Saprolegnia aenigmatica* Sand.-Sierra & Diég.-Urib., PLoS ONE 10:9. 2015. .

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water, surface sediment and submerged leaves of freshwater bodies samples baited with *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4380, CCIBt4406, CCIBt4417. Comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2022).

32. *Saprolegnia atlantica* Pires-Zottar., Rocha S.C.O. & A.L. Jesus, Mycol. Progr. 21 (3, n° 41): 6. 2022.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water of freshwater bodies and soil samples baited with *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4335, CCIBt4351. Description, comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2022).

33. *Saprolegnia blelhamensis* (M.W. Dick) Milko, Mikol. Fitopatol. 13: 290. 1979.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from soil samples baited with *Sorghum* sp seeds, 25 Oct 2017, CCIBt4354, CCIBt4423. Description, comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2022).

34. *Saprolegnia glomerata* (Tiesenh.) A. Lund, Mém. Acad. Roy. Sci. Lett. Sec. Sci 5: 14. 1934.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and surface sediment of freshwater bodies samples baited with *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4352, CCIBt4353, CCIBt4402, CCIBt4549. Description, comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2022).

35. *Saprolegnia milanezii* Pires-Zottar. & S.C.O. Rocha, Phytotaxa 270: 289. 2016.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from water of freshwater bodies samples baited with *Sorghum* sp seeds, 27 Jun 2017, CCIBt4379. Comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2022).

VERRUCALVACEAE

36. *Aphanomyces brasiliensis* Pires-Zottar. & A.L. Jesus, Phytotaxa 415: 214. 2019.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°34'21.1"S, 47°14'53.0"W, from soil samples baited with snake skin 25 Oct 2017,

CCIBT4359, CCIBt4545. GenBank – SSU: OK235679. Description, comments, illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2019).

37. *Aphanomyces helicoides* Minden, Krypt. Fl. Branderburg (Lepizig) 5: 559. 1912 (1915).

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and surface sediment of freshwater bodies samples baited with onion skin, shrimp exoskeleton, snake skin and *Sorghum* sp seeds, 27 Jun 2017 and 25 Oct 2017, CCIBt4331. GenBank – SSU: OK235681. Illustration and sequence code of ITS rDNA region are available in Pires-Zottarelli *et al.* (2019).

38. *Aphanomyces raphani* J.B. Kendr., Phytopathology 17: 43. 1927.

Specimens examined: BRAZIL. SÃO PAULO: Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°33'11.1"S, 47°13'34.5"W, from water of freshwater bodies samples baited with corn leaves, 27 Jun 2017, CCIBt4409. GenBank – SSU: OK235682 Illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2019).

39. *Aphanomyces stellatus* de Bary, J. Wiss. Bot. 2: 178. 1860.

Specimens examined: BRAZIL. SÃO PAULO, Peruíbe and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water, surface sediment and submerged leaves of freshwater bodies and soil samples baited with snake skin, 24 Aug 2016, 27 Jun 2017 and 25 Oct 2017, CCIBt4305. GenBank – SSU: OK235682. Illustration and sequences codes of ITS and LSU rDNA regions are available in Pires-Zottarelli *et al.* (2019).

40. *Phragmosporangium uniseriatum* R.L. Seym., Mycotaxon 92: 8. 2005.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo, Peruíbe and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W ; 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water and submerged leaves of freshwater bodies and soil samples baited with snake skin, shrimp exoeskeleton and *Sorghum* sp seeds, 24 Aug 2016, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4334. GenBank – ITS: ON860616, LSU: ON860626.

Identification: This is the unique species of the genus. *Phragmosporangium uniseriatum* presents abundant production of the zoosporangia and oogonia, oogonial wall with inner surface smooth and outer surface smooth, irregular or papillated; antheridia androgynous and monoclinal and centric and subcentric oospores (Johnson *et al.* 2002, Pires-Zottarelli *et al.* 2007). The morphological characteristics of our specimens are in agreement with the descriptions of these authors, with the ITS rDNA sequences corroborating this identification. It is a new citation for the studied area. Illustration of the species in Pires-Zottarelli *et al.* (2007).

41. *Plectospira myriandra* Drechsler, J. Agric. Res. 34: 295. 1927.

Specimens examined: BRAZIL. SÃO PAULO, Pedro de Toledo, Peruíbe and Iguape, Mosaico de Unidades de Conservação Juréia-Itatins, 24°23'51.5"S, 47°07'14.4"W ; 24°21'59.1"S, 47°18'32.7"W; 24°33'11.1"S, 47°13'34.5"W and 24°34'21.1"S, 47°14'53.0"W, respectively, from water, surface sediment and submerged leaves of freshwater bodies and soil samples baited with onion skin and corn leaves, 24 Aug 2016, 14 Feb 2017, 27 Jun 2017 and 25 Oct 2017, CCIBt4342. GenBank – ITS: ON860618.

Identification: Abundant production of the oogonia, yellowish subcentric oospores, as well as declinal and monoclinal antheridia are important characteristics of this species (Johnson *et al.* 2002, Miranda & Pires-Zottarelli 2012, Jerônimo *et al.* 2017). The morphological characteristics of our specimens are in agreement with the descriptions of these authors. Molecular data corroborated the morphological identification. It is a new citation for the studied area. Illustration of the species in Jerônimo *et al.* (2017).

Discussion

This study allowed us to analyze the occurrence and distribution of the heterotrophic straminipiles of a preserved Brazilian Atlantic Rainforest area. All the identified taxa were isolated from freshwater and soil samples using baits as substrate. The majority of the species was found as saprotrophs, however some of them with pathogenic potential in plants and animals (Plaats-Niterink 1981, Sati 1991, Sandoval-Sierra & Diéguas-Uribeondo 2015, Carraschi *et al.* 2018, Ferreira *et al.* 2021, Dima *et al.* 2021, and others), with several of them with genic regions made available for the first time in GenBank. Many of them have already mentioned for Brazil (Gomes & Pires-Zottarelli 2006, Milanez *et al.* 2007, Mendes *et al.* 1998, Miranda & Pires-Zottarelli 2008, 2012, Nascimento & Pires-Zottarelli 2012, Jesus *et al.* 2013,

Rocha *et al.* 2016a, b, 2018, Jesus *et al.* 2016a, Gonçalves *et al.* 2016a, b), however we contributed with new taxa for the science and new occurrences for the studied area, São Paulo State or Brazil.

Aphanomyces stellatus, *Phragmosporangium uniseriatum* and *Plectospira myriandra* were the most frequent taxa in agreement with other studies in Atlantic Rainforest areas (Pires-Zottarelli *et al.* 2007, Miranda & Pires-Zottarelli 2012), together with *Achlya* sp1 (possibly a new taxon for science).

The calculated species richness and community structure evidenced a diverse microbiota, with few dominant taxa. Although a higher number of species and occurrences were obtained in the soil, there was no statistical difference among the samples. These results are in agreement with Nascimento *et al.* 2011, 2012; Jerônimo *et al.* 2015.

The novelties in relation to this study corroborate the importance of diversity studies, especially in Atlantic Rainforest, a rich Brazilian biome understudied in terms of heterotrophic zoosporic organisms. The obtained sequences enrich the genome database of these organisms.

Acknowledgements

We are indebted to Instituto Florestal, for the permission given to collect samples at “Mosaico de Unidades de Conservação Juréia-Itatins” and to Otto Hartung, for his valuable help during sampling. We also wish to thank Fundação de Amparo à pesquisa do Estado de São Paulo (FAPESP), for the financial support (Process N°. 2016/11146-4), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), for the grant (Process N° 304091/2018-9) given to C.L.A. Pires-Zottarelli.

Conflict of interest

The authors declare no competing interests.

Authors' Contributions

Carmen Lidia Amorim Pires-Zottarelli and Ana Lucia de Jesus: collected freshwater and soil samples, baited them and identified the species using morphological and/or molecular data.

Carmen Lidia Amorim Pires-Zottarelli: wrote the first draft of the manuscript

Ana Lucia de Jesus, Débora Rodrigues da Silva Colombo, Marcela Castilho Boro and Poliana de Oliveira Ventura: help in the final version of the manuscript.

Débora Rodrigues da Silva Colombo and Poliana de Oliveira Ventura: helped us with morphological and/or molecular analysis.

Marcela Castilho Boro: preserved and included the cultures in CCIBt, as well as prepared the plates.

Literature cited

- Adl S.M., Bass D., Lane C.E., Luke, J., Schoch, C.L., Smirnov, A., Agatha, S., Berney, C., Brown, M.W., Burki, F., Ardenas, P.C., Cepi, I., Chistyakova, L., del Campo, J., Dunthorn, M., Edvardsen, B., Eglit, Y., Guillou, L., Hampl, V., Heiss, A.A., Hoppenrath, M., James, T.Y., Karnkowska, A., Karpov, S., Kim, E., Kolisko, M., Kudryavtsev, A., Lahr, D.J.G., Lara, E., Gall, L.L., Lynss, D.H., Mann, D.G., Massana, R., Mitchell, E.A.D., Morrow, C., Park, J.S., Pawlowski, J.W., Powell, M.J., Richter, D.J., Rueckert, S., Shadwick, L., Shimano, S., Spiegel, F.W., Torruella, G., Youssef, N., Zlatogursky, V. & Zhang, Q.** 2019. Revisions to the classification, nomenclature and diversity of eukaryotes. *Journal of Eukaryotic Microbiology* 66: 4-119.
- Beakes, G.W. & Thines, M.** 2017. Hyphochytriomycota and Oomycota. *In: Archibald J, Simpson A, Slamovits C (eds.). Handbook of the Protists.* Springer International Publishing, Cham, pp. 435-505.
- Beakes, G.W., Glockling L.S. & Sekimoto, S.** 2012. The evolutionary phylogeny of the oomycete “fungi”. *Protoplasma* 249: 3-19.
- Beakes, G.W., Honda D. & Thines M.** 2014. Systematics of the Straminipila: Labyrinthulomycota, Hyphochytriomycota, and Oomycota. *In: McLaughlin DJ & Spatafora JW (eds.). The Mycota VIII Part A.* Springer-Verlag, Berlin, pp. 39-97.
- Beneke, E.S. & Rogers, A.L.** 1962. Aquatic Phycomycetes isolated in the States of Minas Gerais, São Paulo and Paraná, Brazil. *Rickia* 1: 181-193.
- Beneke, E.S. & Rogers, A.L.** 1970. Aquatic fungi of Parque Nacional do Itatiaia in the State of Rio de Janeiro. *Rickia* 5: 51-64.
- Carraschi, S.P., Cruz, C. & Ranzani-Paiva, M.J.T.** 2018. Isolation of *Saprolegnia aenigmatica* oomycetes and protocol for experimental infection in pacu (*Piaractus mesopotamicus*). *Acta Scientiarum Biological Science* 33: 324-327.
- Da Paixão, S.C.O.** 2021. Estudo morfológico e molecular de espécies do filo Oomycota isoladas de cultivo de truta arco-íris (*Onchorhynchus mykiss* Walbaum) em Campos do Jordão, estado de São Paulo, Brasil. Tese de Doutorado, Instituto de Botânica, São Paulo.

- Dick, M.W.** 2001. Straminipilous fungi: systematics of the Peronosporomycetes, including accounts of the marine straminipilous protists, the plasmodiophorids, and similar organisms. Kluwer Academic Publishers, Dordrecht.
- Dima, B., Brandrud, T.E., Corriol, G., Janse, G.M., Jordal, J.B., Khalid, A.N., Larsoon, E., Loras, J., Morozova, O.V., Naseer, A., Noordeloos, M.E., Rossi, W., Santamaria, S., Sarwar, S., Sesli, E., Usman, M., Afshann, N.S., Ahmad, I., Banerjee, A., Banerjee, K., Bendiksen, E., Coloambo, D.R.S., De Kesel, A., Dovana, F., Ferisin, G., Hussain, S., Islam, S., Jesus, A.L., Kaygusuz, O., Greilhuber, I.K., Mahammad, S., Mishra, D.K., Nath, P.S., da Paixão, S.C.O., Panja, B., Papp, V., Pires-Zottarelli, C.L.A., Radnóti, A., Rana, D., Saha, R., Türkekul, I & Haelewaters, D.** 2021. Fungal Systematics and Evolution: FUSE 7. *Sydowia* 76: 271-340.
- Ferreira, K.S., Tachibana, L., Colombo, D.R.S., da Paixão, S.C.O., Ferreira, C.M. & Badaró-Pedroso, C.** 2021. Toxic effects in *Aphanomyces brasiliensis* and zebrafish embryos caused by oomycetocides. *Diseases of Aquatic Organisms* 144: 75-87.
- Gomes, A.L. & Pires-Zottarelli, C.L.A.** 2006. Diversidade de Oomycota da Reserva Biológica de Paranapiacaba, Santo André, SP: primeiras citações para o Brasil. *Revista Brasileira de Botânica* 29: 569-577.
- Gonçalves, D.R., Jesus, A.L. & Pires-Zottarelli, C.L.A.** 2016a. *Pythium* and *Phytophythium* species associated with hydroponically grown crops around the city of São Paulo, Brazil. *Tropical Plant Pathology* 41: 397-405.
- Gonçalves, D.R., Jesus, A.L., Rocha, S.C.O., Marano, A.V. & Pires-Zottarelli, C.L.A.** 2016b. New records of *Pythium* (Oomycetes, Straminipila) for South America based on morphological and molecular data. *Nova Hedwigia* 103: 1-12.
- Hammer, Ø., Harper, D.A.T. & Ryan, P.D.** 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4: 1-9.
- Huang, J.H., Chen, C.Y., Lin, Y.S., Ann, P.J., Huang, H.C. & Chung, W.H.** 2013. Six new species of *Pythiogeton* in Taiwan, with an account of the molecular phylogeny of this genus. *Mycoscience* 54: 130-147.
- Huneycutt, M.B.** 1948. Keratinophilic Phycomycetes. I. A new genus of Saprolegniaceae. *Journal of the Elisha Mitchell Scientific Society* 64: 277-285.
- Huneycutt, M.B.** 1952. A new water mold on keratinized materials. *Journal of the Elisha Mitchell Scientific Society* 68: 109-112.
- Jerônimo, G.H., Jesus, A.L., Marano, A.V., James, T.Y., de Souza, J.I., Rocha, S.C.O., Pires-Zottarelli, C.L.A.** 2015. Diversidade de Blastocladiomycota e Chytridiomycota do Parque Estadual da Ilha do Cardoso, Cananéia, SP. *Brasil. Hoehnea* 42: 135-163.

- Jerônimo, G.H., Jesus, A.L., Rocha, S.C.O., Gonçalves, D.R. & Pires-Zottarelli, C.L.A.** 2017. New insights into *Plectospira* genus (oomycetes, Straminipila): morphological and molecular analyses. *Phytotaxa* 307: 191-198.
- Jesus, A.L., Gonçalves, D.R., Rocha, S.C.O., Marano, A.V., Jerônimo, G.H., de Souza, J.I., Boro, M.C. & Pires-Zottarelli, C.L.A.** 2016a. Morphological and phylogenetic analyses of three *Phytopythium* species (Peronosporales, Oomycota) from Brazil. *Cryptogamie, Mycologie* 37: 1-12.
- Jesus, A.L., Marano, A.V., Schoenlein-Crusius, I.H. & Pires-Zottarelli, C.L.A.** 2013. Diversidade de organismos zoospóricos do córrego Pirarungaua, Parque Estadual das Fontes do Ipiranga, São Paulo, Brasil: novas citações. *Hoehnea* 40: 167-180.
- Jesus, A.L., Marano, A.V., de Souza, J.I., James, T.Y., Jerônimo, G.H., Rocha, S.C.O., Gonçalves, D.R., Boro, M.C. & Pires-Zottarelli, C.L.A.** 2015. *Achlya catenulata*, a new Saprolegniales (Oomycetes, Straminipila) from Brazilian mangrove swamp. *Phytotaxa* 212: 221-228.
- Jesus, A.L., Marano, A.V., Jerônimo, G.H., de Souza, J.I., Leão, E.M. & Pires-Zottarelli, C.L.A.** 2016b. The genus *Halophytophthora* (Peronosporales, Straminipila) in Brazil: first descriptions of species. *Brazilian Journal of Botany* 39: 729-739.
- Johnson Jr., T.W.** 1956. The genus *Achlya*: morphology and taxonomy. University of Michigan Press, Michigan.
- Johnson Jr., T.W., Seymour, R.L. & Padgett, D.E.** 2002. Biology and systematics of Saprolegniaceae. Available from <http://dl.uncw.edu/digilib/Biology/Fungi/Taxonomy%20and%20Systematics/Padgett%20Book/Preface.pdf> (access in 28-I-2019)
- Karling, J.S.** 1939. A new fungus with anteriorly uniciliate zoospores: *Hyphochytrium catenoides*. *American Journal of Botany* 44: 512-519.
- Karling, J.S.** 1947. *Lagenidium humanum*, a saprophyte isolated on dead human skin. *Mycologia* 39: 224-230.
- Karling, J.S.** 1977. *Chytridiomycetarum Iconographia*. Lubrecht & Cramer, Vaduz.
- Karling, J.S.** 1987. Two new species of *Leptolegniella* and other zoosporic fungi in Thailand soils. *Nova Hedwigia* 45: 433-443.
- Lyra, N.P. & Milanez, A.I.** 1974. Notas para o levantamento dos ficomicetos aquáticos do Estado de São Paulo. Recife, Instituto de Micologia da Universidade Federal de Pernambuco, Publicação 698: 1-27.
- Marano, A.V., Barrera, M.D., Steciow, M.M., Donadelli, J.L. & Saparrat, M.C.N.** 2008. Frequency, abundance and distribution of zoosporic organisms from Las Cañas stream (Buenos Aires, Argentina). *Mycologia* 100: 691-700.
- Marano, A.V., Gleason, F.H., Rocha, S.C.O., Pires-Zottarelli, C.L.A. & de Souza, J.I.** 2017. Crown Oomycetes Have Evolved as Effective Plant and Animal Parasites. *In*: Dighton J, White F (eds.), The

Fungal Community Its Organization and Role in the Ecosystem. Fourth Edition. CRC press, Taylor and Francis Group, New York, pp. 257-272.

Mendes, M.A.S., Silva, V.L., Dianese, J.C., Ferreira, M.A.S.V., Santos, C.E.N., Gomes Neto, E. & Urben, A.F. (eds.). 1998. Fungos em plantas no Brasil. Embrapa-SPI/Embrapa-Cenargen.

Milanez, A.I. 1989. Fungos de águas continentais. *In*: Fidalgo O, Bononi VL (coords.) Técnicas de coleta, preservação e herborização de material botânico, Série documentos, São Paulo, pp. 17-20.

Milanez, A.I., Pires-Zottarelli, C.L.A. & Gomes, A.L. 2007. Brazilian zoosporic fungi, São Paulo.

Miranda, M.L. & Pires-Zottarelli, C.L.A. 2008. O gênero *Pythium* no Parque Estadual da Serra da Cantareira, estado de São Paulo, SP. Hoehnea 35: 281-288.

Miranda, M.L. & Pires-Zottarelli, C.L.A. 2012. Oomicetos do Parque Estadual da Serra da Cantareira, São Paulo, SP, Brasil. Hoehnea 39: 95-112.

Moreira, C.G. 2006. Avaliação da diversidade e biomassa de fungos associados a folhas em decomposição de *Tibouchina pulchra* Cogn. submersas em reservatórios do Parque Estadual das Fontes do Ipiranga (PEFI), São Paulo, SP. Dissertação de Mestrado, Instituto de Botânica, São Paulo.

Nascimento, C.A. & Pires-Zottarelli, C.L.A. 2012. Diversidade de fungos zoospóricos da Reserva Biológica de Mogi Guaçu, estado de São Paulo, Brasil. Rodriguésia 63: 587-611.

Nascimento, C.A., Gomes, E.P.C. & Pires-Zottarelli, C.L.A. 2011. Occurrence and distribution of zoosporic organisms in water bodies from Brazilian Cerrado areas. Mycologia 103: 261-272.

Nascimento, C.A., Gomes, E.P.C., de Souza, J.I. & Pires-Zottarelli, C.L.A. 2012. Zoosporic true fungi and heterotrophic straminipiles assemblages from soil of Brazilian Cerrado areas. Fungal Ecology 5: 114-123.

Pires-Zottarelli, C.L.A. & Rocha, M. 2007. Novas citações de Chytridiomycota e Oomycota para o Parque Estadual das Fontes do Ipiranga (PEFI), SP, Brasil. Acta Botanica Brasílica 21: 125-136.

Pires-Zottarelli, C.L.A., Colombo, D.R.S., da Paixão, S.C.O., Ventura, P.O., Boro, M.C. & Jesus, A.L. 2019. *Aphanomyces brasiliensis* sp. nov. (Verrucalvaceae, Saprolegniales): a new species from Brazilian Atlantic Rainforest areas. Phytotaxa 415: 208-216.

Pires-Zottarelli, C.L.A., da Paixão, S.C.O., Colombo, D.R.S., Boro, M.C. & Jesus, A.L. 2022. *Saprolegnia atlantica* sp. nov. (Oomycota, Saprolegniaceae) from Brazil, and new synonymizations and epitypifications in the genus *Saprolegnia*. Mycological Progress 21: 41.

Pires-Zottarelli, C.L.A., Gomes, A.L., Oliveira, J. M. & Milanez, A.I. 2007. *Phragmosporangium uniseriatum* in Brazil. Mycotaxon 102: 179-182.

Pires-Zottarelli, C.L.A., Milanez, A.I. & Schoenlein-Crusius, I.H. 1995. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Fungos, 3: Peronosporales. Hoehnea 22: 125-133.

- Pires-Zottarelli, C.L.A., Milanez, A.I., Schoenlein-Crusius, I.H. & Lohmann, L.G.** 1996. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP. Fungos, 4: Saprolegniales. Hoehnea 23: 39-66.
- Plaats-Niterink, A.J. van der.** 1981. Monograph of the genus *Pythium*. Studies in Mycology 21: 1-242.
- Riethmüller, A., Voglmayr, H., Göker, M., Weiß, M. & Oberwinler, F.** 2002. Phylogenetic relationships of the downy mildews (Peronosporales) and related groups based on nuclear large subunit ribosomal DNA sequences. Mycologia 94: 834-849.
- Robideau, G.P., De Cock, A.W.A.M., Coffey, M.D., Hermann, V., Boruwer, H., Bala, K., Chitty, D.W., Désaulniers, N., Eggertson, Q.A., Gachon, C.M.M., Hu, C., Küpper, F.C., Rintoul, T.L., Sarhan, E., Verstappen, E.C.P., Zang, Y., Bonants, P.J.M., Ristaino, J.B. & Lévesque, A.** 2011. DNA barcoding of oomycetes with cytochrome c oxidase subunit I and internal transcribed spacer. Molecular Ecology Resources 11: 1002-1011.
- Rocha, J.R.S.** 2002 Fungos zoospóricos em área de cerrado no Parque Nacional de Sete Cidades, Piauí, Brasil. Tese de Doutorado, Universidade de São Paulo, São Paulo.
- Rocha, J.R., Rodrigues, E.P., Pereira da Silva, H.S.V., Alves de Sousa, L.M. & Skally, B.** 2010. Distribuição geográfica de *Aphanodictyon papillatum* Huneycutt ex Dick (Saprolegniales) no Brasil. Acta Botanica Malacitana 35: 171-176.
- Rocha, S.C.O., Jerônimo, G.H., Marano, A.V., de Souza, J.I., Jesus, A.L. & Pires-Zottarelli, C.L.A.** 2016a. Oomycota from “Parque Estadual da Ilha do Cardoso” (PEIC): first records for São Paulo State and Brazil. Cryptogamie, Mycologie 37: 177-191.
- Rocha, S.C.O., Lopez-Lastra, C.C., Marano, A.V., de Souza, J.I., Ruéda-Paramo, M.E. & Pires-Zottarelli, C.L.A.** 2018. New phylogenetic insights into Saprolegniales (Oomycota, Straminipila) based upon studies of specimens from Brazil and Argentina. Mycological Progress 17: 691-700.
- Rocha, S.C.O., Sandoval-Sierra, J.V., Diéguez-Uribeondo, J., Gonçalves, D.R., Jerônimo, G.H., Jesus, A.L., Marano, A.V. & Pires-Zottarelli, C.L.A.** 2016b. *Saprolegnia milanezii* sp. nov., a new species of Saprolegniales (Oomycota, Straminipila) from Brazil. Phytotaxa 270: 286-294.
- Rogers, A.L., Milanez, A.I. & Beneke, E.S.** 1970. Additional aquatic fungi from São Paulo State. Rickia 5: 93-110.
- Sandoval-Sierra, J.V. & Diéguez-Uribeondo, J.** 2015. A comprehensive protocol for improving the description of saprolegniales (Oomycota): two practical examples (*Saprolegnia aenigmatica* sp. nov. and *Saprolegnia racemosa* sp. nov.) PLoS ONE 10, e0132999.
- Sati, S.C.** 1991. Aquatic fungi parasitic on temperate fishes of Kumaun Himalaya, India. Mycoses 34: 437-441.
- Schmitz, A. & Riesner, D.** 2006. Purification of nucleic acids by selective precipitation with polyethylene glycol 6000. Analytical Biochemistry 354: 311-313.

- Schoenlein-Crusius, I.H. & Milanez, A.I.** 1998. Fungos zoospóricos (Mastigomycotina) da mata atlântica da Reserva Biológica do Alto da Serra de Paranapiacaba, município de Santo André, SP. *Revista Brasileira de Botânica* 2: 177-181.
- Schoenlein-Crusius, I.H., Milanez, A.I., Trufem, S.F.B., Pires-Zottarelli, C.L.A., Grandi, R.A.P., Santos, M.A. & Giustra, K.C.** 2006. Microscopic fungi in the Atlantic rainforest in Cubatão, São Paulo, Brazil. *Brazilian Journal of Microbiology* 37: 244-252.
- Scott, W.W.** 1961. A revision of the genus *Aphanomyces*. Technical Bulletin Virginia Agricultural Experiment Station 151: 1-95.
- Seymour, R.L.** 1970. The genus *Saprolegnia*. *Nova Hedwigia* 19: 1-24.
- Souza, C.R.G. & Souza, A.P.** 2004. Geologia e geomorfologia da área da Estação Ecológica Juréia-Itatins. *In: Marques OAV, Duleba W (eds.) Estação Ecológica Juréia-Itatins Ambiente Físico, Flora e Fauna. Ribeirão Preto: Holos, pp. 16-33.*
- Sparrow Jr., F.K.** 1960. Aquatic Phycomycetes. 2 ed. University of Michigan Press, Ann Arbor.
- Spencer, M.A., Vick, M.C. & Dick, M.W.** 2002. Revision of *Aplanopsis*, *Pythiopsis*, and 'subcentric' *Achlya* species (Saprolegniaceae) using 18S rDNA and morphological data. *Mycological Research* 106: 549-560.
- White, T.J., Bruns, T., Lee, S. & Taylor, J.** 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds.) PCR protocols: A guide to the methods and applications, Academic Press, New York: pp. 315-322.*
- Zak, J.C. & Willig, M.R.** 2004. Fungal biodiversity patterns. *In: Mueller GM, Bills GF, Foster MS (eds.) Biodiversity of Fungi: inventory and monitoring methods, Elsevier Academic Press, USA, pp. 59-75.*

Editora Associada: Elaine Malosso

Submissão: 05/08/2022

Aceito: 09/02/2023

Table 1. Straminipiles from soil and water, submerged leaves and surface sediment from freshwater bodies at “Mosaico de Unidades de Conservação Juréia-Itatins” collected from August/2016 to October/2017. TF%: Total Frequency. TA: Total Abundance.

Taxa/ Samples	Water	Submerged leaves	Sediment	Soil	TF%	TA
STRAMINIPILA KINGDOM						
HYPHOCHYTRIOMYCOTA						
HYPHOCHYTRIALES						
Hyphochytriaceae						
<i>Hyphochytrium catenoides</i>				1	6.25	1
OOMYCOTA						
Leptolegniellaceae <i>incertae sedis</i>						
<i>Aphanodictyon papillatum</i>				1	6.25	1
<i>Leptolegniella keratinophila</i>		1		5	37.50	6
<i>Leptolegniella exogena</i>				1	6.25	1
<i>Leptolegniella</i> sp 1				3	12.50	3
PERONOSPORALES <i>s. l.</i>						
Pythiaceae <i>sensu lato</i>						
<i>Myzocytiopsis humana</i>		3	2	4	75	9
<i>Pythiogeton abundans</i>	3		1		12.50	4
<i>Pythium acanthicum</i>			1	4	50	5
<i>Pythium graminicola</i>			2		12.50	2
<i>Pythium periplocum</i>	1	2			37.50	
<i>Pythium splendens</i>			1		6.25	1
Peronosporaceae <i>s. l.</i>						
<i>Phytophthora undulata</i>		1			6.25	1
<i>Phytopythium chamaehyphon</i>	2	1	4		37.50	7
<i>Phytopythium palingenes</i>	4	3		2	18.75	9
<i>Phytopythium vexans</i>				4	12.50	4
SAPROLEGNIALES						
Saprolegniaceae <i>s. l.</i>						
<i>Achlya catenulata</i>	2	3			25	5
<i>Achlya flagellate</i>	2		2	4	56.25	7
<i>Achlya klebsiana</i>	3	3	2		37.50	8
<i>Achlya diffusa</i>	2	4	1		18.75	7
<i>Achlya orion</i>	4			4	37.50	8
<i>Achlya proliferoides</i>	2				12.50	2
<i>Achlya truncatiformis</i>				1	6.25	1
<i>Achlya</i> sp 1	6	6	7	7	75	26
<i>Aplanopsis terrestris</i>				1	6.25	1
<i>Brevilegnia longicaulis</i>			1	2	12.50	3
<i>Brevilegnia</i> sp 1				2	12.50	2
<i>Dictyuchus pseudodictyon</i>		3	1		37.50	4

<i>Leptolegnia eccentrica</i>				3	12.50	1
<i>Newbya androgyna</i>	1			2	25	3
<i>Pythiopsis irregularis</i>				3	12.50	3
<i>Saprolegnia aenigmatica</i>	3	1	1		37.50	7
<i>Saprolegnia belhamensis</i>				2	6.25	2
<i>Saprolegnia glomerata</i>	2	1	3		37.50	6
<i>Saprolegnia milanezii</i>	1				6.25	1
<i>Saprolegnia atlantica</i>	3			1	25	4
Verrucalvaceae						
<i>Aphanomyces brasiliensis</i>			1	1	25	2
<i>Aphanomyces helicoides</i>	3	1	1		37.50	5
<i>Aphanomyces raphanin</i>	1				6.25	1
<i>Aphanomyces stellatus</i>	1	3	5	1	75	10
<i>Phragmosporangium uniseriatum</i>	1	1		12	75	14
<i>Plectospira myriandra</i>	6	3	3	4	100	16
Total of species	21	17	18	25		
Total of occurrences	53	40	39	77		209

Table 2. Species richness (S), evenness (E), Shannon diversity (H') and Simpson dominance (D) indices in each of the samples collected. The same letters indicate that the values do not differ significantly ($P \leq 0.05$).

Index	Water	Submerged leaves	Sediment	Soil
S	21a	17a	18a	25a
E	0.85a	0.85a	0.79a	0.77a
H'	2.89a	2.67a	2.65a	2.96a
D	0.06a	0.08a	0.09a	0.06a




Figure 1. Map of the “Mosaico de Unidades de Conservação Juréia-Itatins”, São Paulo State, Brazil, showing the municipalities sampled (adapted from Souza and Souza 2004).




Figure 2. a-b. *Hyphochytrium catenoides* Karling. a. Young thallus on snake skin. b. Polycentric thallus on snake skin with intercalary and terminal swellings and zoosporangia connected by hyphae. c-d. *Aphanodictyon papillatum* Huneycutt ex M.W. Dick. c. Zoosporangia in snake skin. d. Papillate oogonium with thick oospores on snake skin. e. *Pythiogeton abundans* J.H. Huang, Chi Y. Chen & Yi S. Lin. Zoosporangium. f-g. *Pythium acanthicum* Drechsler. f. Intercalary ornamented oogonium with oospore and monoclinous antheridium. g. Details of the oogonia with conical projections. H. *Pythium splendens* Hans Braun. Globose hyphal swellings with dark granular content in the mycelium. Scale bars = 10µm




Figure 3. a-b. *Phytophthora undulata* (H.E. Petersen) M.W. Dick. a. Zoosporangium with internal proliferation. b. Chlamydospore. c-d. *Achlya diffusa* J.V. Harv. Ex T.W. Johnson. C. “Balloon-like” hyphal swelling. d. Oogonia with eccentric oospores and declinuous antheridia. e-f. *Achlya truncatiformis* M.W. Dick & Mark A. Spencer. E. Achlyoid discharge of the zoosporangia. f. Oogonia with truncate oogonial papillae and eccentric oospores. g-h. *Pythiopsis irregularis* R.L. Seym. g. Empty zoosporangia and initial developmental of oogonium and antheridia. h. Ornamented oogonium with subcentric oospores. Scale bars = 10 μ m

CARTA DE AUTORIZAÇÃO DE PUBLICAÇÃO NO PORTAL DE PREPRINTS DO SCIELO

Ao Comitê Editorial de HOEHNEA

Declaro, em meu próprio nome e nos dos demais Autores, que concordo com a publicação do Artigo Aceito pelo Corpo Editorial de Hoehnea, intitulado “Occurrence and distribution of the heterotrophic straminipiles from Brazilian Atlantic Rainforest areas” de autoria de Carmen Lidia Amorim Pires-Zottarelli, Ana Lucia de Jesus, Débora Rodrigues da Silva Colombo, Poliana de Oliveira Ventura e Marcela CastilhoBoro, no Portal de Preprints do SciELO Brasil (<https://preprints.scielo.org/index.php/scielo/preprints/section/biological>).

Declaro, ainda, que o referido artigo é original, sendo que o conteúdo não foi ou não está sendo considerado para publicação em outro periódico, quer seja no formato impresso e/ou eletrônico.

São Paulo, 13 de fevereiro de 2023.


Carmen Lidia Amorim Pires Zottarelli

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.