

Publication status: Not informed by the submitting author

Biological, social, and healthcare factors associated with death due to influenza A(H1N1) during the 2009 epidemic in Brazil

Jose Ueleses Braga, Ana Freitas Ribeiro

<https://doi.org/10.1590/SciELOPreprints.6696>

Submitted on: 2023-09-11

Posted on: 2023-09-15 (version 1)

(YYYY-MM-DD)

Biological, social, and healthcare factors associated with death due to influenza A(H1N1) during the 2009 epidemic in Brazil

Jose Ueleres Braga

Instituto de Medicina Social Hesio Cordeiro / UERJ, Departamento de Epidemiologia. Rio de Janeiro – RJ. Brasil. ORCID: <https://orcid.org/0000-0001-5247-007X>

Escola Nacional de Saúde Pública Sergio Arouca / FIOCRUZ, Departamento de Epidemiologia de doenças infecciosas. Rio de Janeiro – RJ. Brasil.

Ana Freitas Ribeiro

Instituto de Infectologia Emílio Ribas - Secretaria de Estado da Saúde de São Paulo - São Paulo – SP. Universidade Municipal de São Caetano do Sul - Faculdade de Medicina - São Paulo - SP. ORCID: <https://orcid.org/0000-0001-9111-4510>

ABSTRACT

Background: In June 2009, the World Health Organization (WHO) raised the global alert level for the A(H1N1)pdm09 influenza pandemic and at that time sustained transmission in Brazil was established. It was urgent to carry out studies that evaluated possible risk factors for death from Influenza A(H1N1) to improve case management strategies to reduce the lethality of the disease. This study aimed to identify risk factors for death from Influenza A(H1N1), including the effectiveness of the vaccine against influenza A(H1N1) concerning mortality. Methods: A case-control of incident cases of influenza A(H1N1) reported in the Epidemiological Information Systems of the states of São Paulo, Paraná, Pará, Amazonas, and Rio Grande do Sul was conducted. Results: 305 participants were included, 70 of them cases and 235 controls, distributed as follows: Amazonas – 9 cases/10 controls, Pará – 22 cases/77 controls, São Paulo – 19 cases/49 controls, Paraná – 10 cases/54 controls, Rio Grande do Sul – 10 cases/45 controls. These participants had a mean age of 30 years, with 33 years among cases and 25 years among controls. There was a predominance of females both among cases and controls. Biological (age), pre-existing diseases (congestive heart failure, respiratory disease, and diabetes mellitus), and care factors (ICU admission) associated with death from Influenza A(H1N1) were identified. Conclusion: The risk factors identified in this investigation allowed subsidizing the elaboration of clinical conducts, but also indicate important aspects for facing “new” influenza epidemics that are likely to occur in our country.

Keywords:

“influenza a virus, h1n1 subtype”, “disease outbreaks”, “risk factors”, “case-control studies”.

1. Introduction

In April 2009, the CDC reported two cases of febrile respiratory illness in children in Southern California with laboratory diagnosis of infection with a virus genetically similar to Swine Influenza A H1N1. The virus contains a gene segment not previously identified in human and swine influenza. The two children had no contact with the pigs. Another six cases have been reported to the California and Texas Departments of Public Health. At the same time, samples from patients from Mexico confirmed the same viral subtype.⁽¹⁾

On April 12, 2009, Mexico recorded an outbreak of acute respiratory disease in Gloria, Veracruz, characterized by a high attack rate (28.5%). In March and April, 47 cases of severe pneumonia with 12 deaths were identified in Mexico City, San Luis de Potosi, and other cities. In four samples, the new viral subtype of influenza A(H1N1) was identified.⁽²⁾

On June 11, 2009, the World Health Organization (WHO) raised the global alert level for the pandemic caused by the new influenza A/H1N1 virus – swine lineage – to phase 6 (Pandemic). The action occurred due to the rapid spread of the virus and not due to the severity of the disease.⁽¹⁾ The vast majority of cases evolved to cure. However, it was expected that the number of cases, hospitalizations, and deaths would increase until the end of the epidemic.⁽³⁾ In August 2010, the World Health Organization declared the end of the pandemic, with approximately 18,500 deaths, moving into the post-pandemic phase, with seasonal transmission of the virus.⁽⁴⁾

Sustained transmission in Brazil was established on July 16, 2009. From that date, only cases of Severe Acute Respiratory Syndrome (SARI) were reported, hospitalized, and treated with oseltamivir. Outpatient cases, if considered risk patients (comorbidities, pregnant women, immunosuppressed) could also receive specific treatment up to 48 hours after the onset of symptoms, after medical evaluation. However, in this case, there was no clinical sample collection for diagnosis and no record in the notifiable health information system (SINAN).⁽⁵⁾

At that time, it was believed to be necessary to maintain monitoring of possible changes in the profile of viral circulation and the worsening of the clinical picture of cases of influenza A(H1N1), due to the concomitant circulation of different subtypes of influenza viruses and new evidence of predictive factors. severity and deaths related to this new disease.⁽⁶⁾

Therefore, carrying out studies that evaluated possible risk factors for death from Influenza A(H1N1) and for death in pregnant women were essential for a better understanding of this epidemic. This study aimed to identify risk factors for death from Influenza A(H1N1), a new viral subtype associated with Severe Acute Respiratory Disease (DRAG), including the effectiveness of the H1N1 vaccine in terms of mortality, based on clinical epidemiological data of cases reported to the Brazilian Epidemiological Surveillance System.

2. Methods

This case-control study included incident cases of influenza A(H1N1) reported in the Epidemiological Information Systems of the states of São Paulo, Paraná, Pará, Amazonas, and Rio Grande do Sul. The study population included children, adolescents, and adults with symptoms compatible with suspected infection by the influenza A(H1N1) virus treated at the Unified Health System.

Individuals included as cases had confirmed influenza A(H1N1) infection with fever, respiratory symptoms, and positive confirmatory test for the presence of the virus and registered as death in the Epidemiological Surveillance System for Influenza – SIVEP-Flu or SIM (Information System on Mortality) that were evaluated in loco, that is, their medical records were evaluated.

A time-paired case-control design was used⁽⁷⁾ by the epidemiological week of hospitalization and H1N1 notification records were drawn from the SIVEP-Flu database,

according to the ratio of four controls for each case.⁽⁸⁾

Firstly, the medical records were evaluated and the questionnaire for the first two selected controls was completed. The interviewer sent the completed questionnaires to the field supervisor, who also evaluated in order to verify that the controls fulfilled the case definition. If they did not, the other randomly selected controls had their medical records evaluated. Two controls were sought per case, and a maximum of four medical records of possible patients eligible for control were evaluated.

The sources of data were, therefore: (a) medical records, (b) data or information obtained from family members through home visits and (c) data from the Influenza Epidemiological Surveillance System – SIVEP-Flu. A standardized form was applied to collect data from medical records and home interviews, both for cases and controls.

The variables studied were related to (i) time: date of onset of symptoms, date of first consultation, date of hospitalization, date of death; (ii) place: place of residence, place of outpatient and hospital care; (iii) person: age, sex, education, self-reported race/color, occupation, underlying diseases, vaccination; (iv) clinical and laboratory characteristics: initial clinical and laboratory evaluation, radiological imaging, use of O₂, presence of comorbidities, pregnancy (when applicable), associated infections, other diagnoses during hospitalization, complications during hospitalization, treatment with oseltamvir (dose, time of use and interval of use), cause of death, data from the death certificate, necropsy report (when available), anatomopathological reports and relevant laboratory results (cultures performed).

The size was calculated based on a 5% alpha error, and 20% beta error and indicated the need for 80 A(H1N1) influenza deaths and 320 controls, but if the association estimate had a magnitude greater than 3.0, fewer participants would be needed.

This study was approved by the Research Ethics Committee of the Institute of Social Medicine of the State University of Rio de Janeiro, on December 4, 2009 (CAEE 0030.1.259.000-09).

3. Results

A total of 305 participants were included, 70 cases and 235 controls. Two federal units of the original project were replaced (Rio de Janeiro and Minas Gerais) by the states of Amazonas and Pará. The number of cases and controls in each federal unit was: Amazonas – 9 cases/10 controls, Pará – 22 cases/77 controls, São Paulo – 19 cases/49 controls, Paraná – 10 cases/54 controls, Rio Grande do Sul – 10 cases/45 controls.

Cases and controls were recruited in these federated units in 2010 and 2011, and most participants had symptom onset in the first year of the study. At that time, the vaccine had already been applied to the population and this would also allow evaluating the effect of this intervention on the risk of death from influenza A(H1N1).

The states of Amazonas and Pará were the ones that included the study participants later because the number of deaths in the country had declined in such a way that it only negatively influenced recruitment in these states.

These participants had a mean age of 30 years, with 33 years among cases and 25 years among controls. There was a predominance of females both among cases and controls. The age group with the highest frequency of participants was 18 to 49 years old (Table 1).

Regarding the clinical risk factors for death from Influenza A(H1N1) between cases and controls, differences in the study for the occurrence of some clinical conditions were also

observed in the scientific literature. Cases most often had the following clinical conditions: obesity, systemic arterial hypertension, cerebrovascular disease, congestive heart failure, diabetes mellitus, and respiratory disease. Pregnancy was more frequent among cases than controls, but this difference was not statistically significant at the 5% level (Table 1).

The most frequent signs and symptoms were fever (94.1%), cough (91.1%), dyspnea (80.3%), and headache (33.8%). Regarding its distribution between cases and controls, there are no important differences regarding the characteristic symptoms of the respiratory disease. However, dyspnea was more frequent in those who died (Table 2).

The treatment that cases and controls underwent was quite different than expected. There was no difference concerning the use of Oseltamivir since this medication was available and used very frequently at that time. However, ICU admission and transfer to another service occurred more among cases when compared to controls. The most observed clinical complications were shock and sepsis, in practically 2/3 of the patients who died. All other complications were observed among the cases (Table 2).

The therapeutic procedures used in cases and controls were also different, being more frequent in those individuals who died. The use of antibiotics was different in the two groups, indicating that secondary infections may have occurred more between cases. We highlight the use of invasive and non-invasive ventilation, anticoagulants, and nasogastric tubes. These findings indicate, on the one hand, that patients who died received intensive treatment, but also that these procedures were available to individuals in the control group (Table 2).

The vaccination status of the participants was different between the study sites. Participants from Paraná were more vaccinated (15.6%) than the other states: Amazonas with 5.3%, Pará with 1%, and Rio Grande do Sul with 6.3%. In the state of São Paulo, no participant had received the vaccine. On average, only 5% of study participants were vaccinated.

Regarding pre-existing diseases, it can be said that they influence the prognosis of

patients, increasing the risk of death from influenza A(H1N1). It is noticed that not only chronic respiratory diseases but systemic diseases are associated with death, such as diabetes mellitus and congestive heart failure. The analysis of the joint influence of these morbid conditions revealed that, although the respiratory situation is important, they are not the greatest risk factors (Table 3). Our study did not detect that morbid conditions characterized by immunosuppression had an increased risk for our outcome.

To answer the research question – what are the risk factors for death from Influenza A(H1N1) in Brazil during the epidemic, we decided to use models that encompass the various dimensions studied, biological characteristics, pre-existing diseases, and assistance hospitals received by patients. The first model included only the first two dimensions, the second model also included service transfer and the third also considered ICU admissions.

In model 1 (Table 4) it is evident that both biological factors such as age and obesity can influence the risk of dying from influenza A(H1N1), but it is also influenced by pre-existing diseases such as congestive heart failure and diabetes mellitus. The results of model 2 indicate that assistance can also independently influence the risk of the studied outcome. Finally, in the model that includes ICU admission, many factors “lose their ability to discriminate the risk of death”, which does not mean that these are not risk factors, but that the risks of these biological and clinical conditions are probably represented on the severity of the critical condition requiring intensive care.

4. Discussion

Our results indicate that biological factors such as age and obesity are related to the risk of dying from influenza A(H1N1). There is also evidence that pre-existing diseases such as congestive heart failure, respiratory disease, and diabetes mellitus are associated with

death from this cause. However, our study reveals that the transfer of hospitalization service and ICU stay are conditions that imply that the worsening of the clinical condition increases the risk of death due to influenza A(H1N1).

The set of signs presented by the participants are typical of respiratory syndrome and the absence of significant difference in the set of signs and symptoms indicates the adequacy of the study selection process about the eligibility criteria. This profile was similar to that observed in other investigations carried out in Brazil.⁽⁹⁾⁽¹⁰⁾

Ribeiro et al. 2015⁽¹⁰⁾ conducted a case-control study with 193 reported and confirmed deaths of patients with influenza A(H1N1) and 386 patients hospitalized during the study period with reported and confirmed influenza A(H1N1) who recovered. The distribution of cases and controls according to demographic variables indicated that the median age of cases was higher than that of controls and the age group from 18 to 59 years old had a higher risk compared to those aged less than 18 years old. The elderly also had a significant risk of dying from influenza A(H1N1).

During the period of the 2009 epidemic, another case-control study was also conducted in Brazil. Yokota et al. 2011⁽⁹⁾ carried out an investigation in 11 hospitals in 4 cities (Passo Fundo, Caxias do Sul, Santa Maria, and Uruguaiana) in Rio Grande do Sul, Brazil. The study included 157 patients, 52 patients who died, and 105 who survived. A total of 136 sought treatment before hospitalization and obesity was the most common underlying medical condition (38%). Among the obese cases, 19 (36%) had other risk factors for influenza complications. Diabetes was the most frequent medical condition (21%) in the obese.

In our investigation, most of the factors studied were not associated with the outcome of death from Influenza A(H1N1), so the results of the statistical analysis of the data included only the factors whose underlying theory or exploratory analysis indicated a possible association. Although obesity was not included in the final model, this condition was associated with death from influenza A(H1N1) in our study.

Sun et al. 2016⁽¹¹⁾ performed a systematic review combining results from 22 articles involving 25,189 laboratory-confirmed patients. Pooled estimates indicated that obesity significantly increased the risk of death from influenza A (H1N1). However, the authors indicate that they found a significant interaction between early antiviral treatment and obesity. After adjusting for early antiviral treatment, the relationship between obesity and poor outcomes disappeared. This situation may also have occurred in our study. A review study carried out by Karlsson et al. (2012) ⁽¹²⁾ showed that obesity is associated with decreased lung function, as well as the development of chronic respiratory conditions, and risk factors for the development of severe influenza-associated infection.

Balaganesakumar et al. 2013⁽¹²⁾ conducted a case-control study in Tamil Nadu, India, with 280 participants, 70 patients who died and 210 patients who recovered, between July and December 2010. In this study, fever (96%) and cough (79%) were the most common symptoms, while 12% of patients had dyspnea and 85% of deaths. This profile was similar to that of our study, with a higher proportion of fever (94.1%) and dyspnea (80.3%) than the other signs. In the Indian study⁽¹²⁾, , the chances of death were higher in diabetics, and obese, who received treatment from private services and treated with corticosteroids.

Martinez et al. 2019⁽¹³⁾ conducted an observational study with patients aged ≥ 18 years from 12 Catalan hospitals, in 2010-2016. A total of 1,726 hospitalized patients were included, 595 (34.5%) were admitted to the ICU, and 224 (13.0%) died. Age groups 65-74 and ≥ 75 years were associated with an increased risk of death for all types and subtypes, especially for influenza B. Immunodeficiency was associated with death for influenza B and subtype A(H1N1). These findings are similar to our study, but immunodeficiency was not in the model.

Mata-Marin 2015⁽¹⁴⁾ conducted a case-control study to determine the risk factors associated with death from influenza A(H1N1), detected by RT-PCR. The study included patients over the age of 18 treated at the Hospital de Infectología, Centro Médico Nacional "La Raza", in Mexico City, between April and November 2009. Cases were patients who died

during hospitalization, while controls received high after getting better. A total of 33 patients met the eligibility criteria. Risk factors associated with mortality were: male gender, delayed medical care >3 days, delayed influenza therapy >3 days, intensive care unit (ICU) admission, and creatinine levels >1.0 mg/dL when admitted to the hospital. After adjusting for a logistic regression model, delay in medical care and ICU stay were the only predictors of mortality.

Peralta et al. 2010⁽¹⁵⁾ studied factors associated with the risk of death in cases of 2009 pandemic (H1N1) virus infection hospitalized in intensive care units (ICU) in Spain. Of the 1,231 cases admitted to the ICU, 271 died. The median age was 40 years, 76.3% of patients had some previous disease, respiratory disease being the most frequent (34.1%), followed by morbid obesity (18.8%). In multivariate analysis, cancer, immunodeficiencies, and morbid obesity were significantly associated with death in adults.

Another Brazilian study⁽¹⁶⁾ addressed severe maternal morbidity due to respiratory diseases and the impact of the 2009 influenza A(H1N1) pandemic in Brazil. The objective of this study was to evaluate the burden of respiratory diseases, considering the time of the pandemic within the scope of the Brazilian Network for the Surveillance of Severe Maternal Morbidity, and the factors associated with the worst maternal outcome. Women with severe complications from respiratory disease identified as suspected or confirmed cases of influenza A(H1N1) or respiratory failure were compared with those with other causes of severe morbidity. A review of suspected influenza A(H1N1) cases classified women as untested, tested positive, and tested negative, comparing their results.

Ribeiro et al. 2018⁽¹⁷⁾ also studied the factors associated with death and described the gestational outcomes in pregnant women with A(H1N1)pdm09 influenza. severe acute respiratory illness (SARI) in the state of São Paulo from June 9 to December 1, 2009. We investigated 48 cases and 185 controls. The results presented in this study indicate that early treatment can prevent unfavorable outcomes for pregnant women and their children and reinforce the need for adequate training of doctors for the clinical management of pregnant women and early administration of antiviral treatment.

No other Brazilian study was located in the scientific literature that had a scope and methodological approach similar to that used in our investigation. The challenge of identifying risk factors for death from H1N1 Influenza includes the need to also consider the medical and hospital care provided to patients.

If, on the one hand, it is worth saying that adequate and timely assistance can prevent deaths, it must also be considered that the most critical patients are those who receive intensive care. Therefore, we questioned whether the transfer of patients to more specialized services and admission to an intensive care unit was associated with the risk of dying from H1N1 influenza. The occurrence of “reverse causality” is plausible if these relationships are considered causal. This study intended to indicate risk markers for death from this disease, even though we use the term risk factor to indicate these studied characteristics. As expected, that is, hypothesized, both transferring services and hospitalization were strongly associated with death from influenza. The strength of the association found for ICU admission is the highest found in our investigation.

Regarding the ability of this study to assess the influence of the participants' vaccination status on the risk of dying from A(H1N1) influenza, it is worth considering that the low proportion of vaccinated individuals in the sample may have had a negative influence. Despite the result indicating protection when calculating the measure of effect adjusted for the other factors, only 5% of the participants were vaccinated and, as already mentioned, the expected effect of the immunization campaign may not yet have been detected. In our study, participants vaccinated for Influenza were not more among the cases (deaths) than non-vaccinated ones. In fact, in March 2010, the federal government started the vaccination campaign. In just three months, using the vaccines purchased and new batches manufactured by the Butantan Institute, Brazil managed to vaccinate 92 million people. The federal government expected to vaccinate 80% of priority groups but managed to reach 88%. The success is due to the mobilization effort and the strategy of implementing vaccination points

in schools, public offices, workplaces, and even public roads.⁽¹⁸⁾

Mahmud et al. 2011⁽¹⁹⁾ performed a population-based case-control study to evaluate the effectiveness of pandemic vaccines against influenza A(H1N1) during the 2009 mass vaccination campaign in Manitoba (Canada). Cases were individuals who tested positive for influenza A by PCR (N=1,435). Controls were individuals with negative results for influenza A and B (N=2,309). Results indicated that the adjuvanted H1N1 vaccine was 86% effective (95% CI 75–93%) in preventing H1N1 infections when vaccination occurred ≥ 14 days before testing. Although it was not possible to study the effectiveness of the vaccine in our study, there seems to be no doubt about the benefits.

This study had limitations such as underreporting of SARS, but as the epidemic progressed, it may have decreased. The quality of hospital record data may also have influenced the results. The strategy used to minimize the quality of data extraction was the careful review of this process by the supervisor.

Our study had to consider the need to address aspects of various dimensions to identify the risk factors for death from influenza A(H1N1) and the three final models reveal the importance of the groups of factors studied. These approaches complement each other in a way because in the absence of a robust theoretical model to study this question, highlighting the importance of these characteristics is only possible with this analytical approach.

The risk factors identified in this investigation allowed for subsidizing the elaboration of clinical conducts, but also indicated important aspects for facing “new” influenza epidemics that are likely to occur in our country.

5. Acknowledgments

We appreciate the help of collaborators Bernardino Claudio de Albuquerque, Geraldine

Madalosso, Letícia Garay Martins, Marilina Assunta Bercini, Suzana Dal-Ri Moreira and Tânia do Socorro Souza Chaves.

6. Financial support

This research was carried out with the financial support of the Convenio 728336/2009 of FOMENTO À PESQUISA EM SAÚDE of Departamento de Ciência e Tecnologia - Ministério da Saúde do Brasil.

7. Conflict of interests

The authors have no affiliation with any organization with a direct or indirect financial interest in the subject matter discussed in the manuscript.

8. Rerefences

1. Centers for Disease Control and Prevention (CDC). Novel influenza A (H1N1) virus infections in three pregnant women - United States, April-May 2009. *MMWR Morb Mortal Wkly Rep* 2009;58(18):497–500.
2. Charatan F. Mexico City returns to normality, but cases of A/H1N1 reach almost 2000 in US. *Bmj* 2009;338:b1950.
3. WHO. Pandemic (H1N1) 2009, weekly update 76. n.d.
4. PAHO, WHO. WHO Declares H1N1 Pandemic Over, Urges Continued Influenza Vigilance. 2010.
5. Ministério da Saúde, Secretaria de Vigilância em Saúde. Informe Epidemiológico, Influenza A H1N1, Situação epidemiológica da nova influenza A (H1N1) no Brasil, até semana epidemiológica 44 de 2009. 2009.
6. Ministério da Saúde. Protocolo de manejo clínico e vigilância epidemiológica da influenza. Brasília: 2009.
7. Wacholder S. Design issues in case-control studies. *Stat Methods Med Res* 1995;4(4):293–309.
8. Schlesselman JJ, Stolley PD. Case-control studies : design, conduct, analysis. New York: Oxford University Press; 1982.
9. Yokota R. Risk Factors for Death from Pandemic (H1N1) 2009, Southern Brazil. *Emerg Infect Dis* 2011.

10. Ribeiro AF, Pellini AC, Kitagawa BY, Marques D, Madalosso G, de Cassia Nogueira Figueira G, et al. Risk factors for death from Influenza A(H1N1)pdm09, State of São Paulo, Brazil, 2009. *PLoS One* 2015;10(3):e0118772.
11. Sun Y, Wang Q, Yang G, Lin C, Zhang Y, Yang P. Weight and prognosis for influenza A(H1N1)pdm09 infection during the pandemic period between 2009 and 2011: a systematic review of observational studies with meta-analysis. *Infect Lond* 2016;48(11–12):813–22.
12. Balaganesakumar SR, Murhekar MV, Swamy KK, Kumar MR, Manickam P, Pandian P. Risk factors associated with death among influenza A (H1N1) patients, Tamil Nadu, India, 2010. *J Postgrad Med* 2013;59(1):9–14.
13. Martínez A, Soldevila N, Romero-Tamarit A, Torner N, Godoy P, Rius C, et al. Risk factors associated with severe outcomes in adult hospitalized patients according to influenza type and subtype. *PLoS One* 2019;14(1):e0210353.
14. Mata-Marín LA, Mata-Marín JA, Vásquez-Mota VC, Arroyo-Anduiza CI, Gaytán-Martínez JE, Manjarrez-Téllez B, et al. Risk factors associated with mortality in patients infected with influenza A/H1N1 in Mexico. *BMC Res Notes* 2015;8(1):432.
15. Santa-Olalla Peralta P, Cortes García M, Limia Sánchez A, Andrés Prado J, Pachón Del Amo I, Sierra Moros MJ. Casos de infección por gripe pandémica (H1N1) 2009 hospitalizados en cuidados intensivos en España: factores asociados a riesgo de muerte, abril 2009-enero 2010. *Rev Esp Salud Pública* 2010;84(5):547–67.
16. Pfitscher LC, Cecatti JG, Pacagnella RC, Haddad SM, Parpinelli MA, Souza JP, et al. Severe maternal morbidity due to respiratory disease and impact of 2009 H1N1 influenza A pandemic in Brazil: results from a national multicenter cross-sectional study. *BMC Infect Dis* 2016;16:220.
17. Ribeiro AF, Pellini ACG, Kitagawa BY, Marques D, Madalosso G, Fred J, et al. Severe influenza A(H1N1)pdm09 in pregnant women and neonatal outcomes, State of Sao Paulo, Brazil, 2009. *PLoS One* 2018;13(3):e0194392.
18. CEE, Centro de Estudos Estrategicos da FIOCRUZ Antonio Ivo de Carvalho. Combate à epidemia de H1N1: um histórico de sucesso. 2021.
19. Mahmud S, Hammond G, Elliott L, Hilderman T, Kurbis C, Caetano P, et al. Effectiveness of the pandemic H1N1 influenza vaccines against laboratory-confirmed H1N1 infections: Population-based case–control study. *Vaccine* 2011;29(45):7975–81.

Table 1 – Distribution of demographic characteristics and risk conditions of participants in the case-control study for risk factors for death due to Influenza A(H1N1)

Characteristic	Cases		Controls		total	p-value
	n	%	n	%		
Sex						
Male	18	18.0	82	82.0	100	0.1509
female	52	25.4	153	74.6	205	
Age						
up to 18 years	15	14.9	86	85.1	101	0.0106
19 to 49 years old	36	25.9	103	74.1	139	
50 years and over	16	37.2	27	62.8	43	
mean age	33.6		25.4			0.0009
Median age	34		23			
Risk conditions						
Pregnant	14	26.9	38	73.1	52	0.0711
quarter						
	1	0	0.0	5	100.0	5
	2	3	15.0	17	85.0	20
	3	10	38.5	16	61.5	26
Puerperium	1	25.0	3	75.0	4	0.0002
Obesity	10	55.6	8	44.4	18	0.1036
Cardiovascular	7	43.8	9	56.3	16	0.0010
Disease						
Hypertension	15	50.0	15	50.0	30	0.1835
Coronary Disease	1	100.0	0	0.0	1	0.1835
Cerebrovascular	2	100.0	0	0.0	2	0.0279
Disease						
Heart Failure	4	80.0	1	20.0	5	0.0084
Diabetes	7	87.5	1	12.5	8	0.0001
Thyroid disease	3	37.5	5	62.5	8	0.6074
Respiratory disease	17	38.6	27	61.4	44	0.0250
Asthma	6	21.4	22	78.6	28	0.9703
COPD	6	46.2	7	53.8	13	0.1011
Tuberculosis	1	50.0	1	50.0	2	0.6119
Cystic fibrosis	0		0		0	
liver disease	2	40.0	3	60.0	5	0.6488
Hemoglobinopathy	0		0		0	
DAI	1	25.0	3	75.0	4	0.7615
Lupus	0	0.0	1	100.0	1	0.6123
Rheumatoid arthritis	0	0.0	2	100.0	2	0.5519
Others DAI	1	50.0	1	50.0	2	0.4921
Kidney disease	1	100.0	0	0.0	1	0.1341
Others diseases	9	21.4	33	78.6	42	0.9645
HIV	1	16.7	5	83.3	6	0.9049
Organ transplantation	0	0.0	1	100.0	1	0.8220
Neoplasm	1	25.0	3	75.0	4	0.9767

Immunosuppressive drug	1	16.7	5	83.3	6	0.7986
---------------------------	---	------	---	------	---	--------

Table 2 – Distribution of signs and symptoms, hospitalization and procedures of participants in the case-control study for risk factors for death due to Influenza A(H1N1)

Característica	Casos		Controles		total	valor de p
	n	%	n	%		
Signs and symptoms						
Fever	62	21.6	225	78.4	287	0.0053
Cough	62	22.3	216	77.7	278	0.2067
Dyspnea	67	27.3	178	72.7	245	0.0004
Headache	10	9.7	93	90.3	103	0.0001
Chill	9	26.5	25	73.5	34	0.2774
Sore throat	10	13.9	62	86.1	72	0.1776
Arthralgia	2	7.1	26	92.9	28	0.0658
Myalgia	16	17.0	78	83.0	94	0.1602
Conjunctivitis	0	0.0	7	100.0	7	0.1760
Runny nose	13	13.8	81	86.2	94	0.0127
Diarrhea	13	34.2	25	65.8	38	0.0611
Vomiting	11	19.0	47	81.0	58	0.3556
Seizures	1	50.0	1	50.0	2	0.2407
Asthenia	10	14.9	57	85.1	67	0.1366
Lack of appetite	7	16.3	36	83.7	43	0.1288
Irritability	1	25.0	3	75.0	4	0.5550
Others	39	24.1	123	75.9	162	0.7930
Hospitalization						
Transfer from another service	36	41.4	51	58.6	87	0.0000
ICU admission	60	64.5	33	35.5	93	0.0000
Use of Oseltamivir	65	23.8	208	76.2	273	0.4240
Complications	59	67.8	28	32.2	87	0.0000
Shock	42	89.4	5	10.6	47	0.0000
Sepsis	45	80.4	11	19.6	56	0.0000
Clotting Disease	8	88.9	1	11.1	9	0.0000
Acute Respiratory Distress Syndrome	36	80.0	9	20.0	45	0.0000
Pleural disease	10	66.7	5	33.3	15	0.0001
Pulmonary hemorrhage	10	90.9	1	9.1	11	0.0000
Other lung diseases	23	59.0	16	41.0	39	0.0000
Heart failure	14	93.3	1	6.7	15	0.0000
Kidney disease	25	86.2	4	13.8	29	0.0000
Procedures						
use of antibiotics	66	25.9	189	74.1	255	0.0137
Oxygen therapy	58	36.5	101	63.5	159	0.0000
Non-invasive ventilation	29	64.4	16	35.6	45	0.0000
Invasive ventilation	53	81.5	12	18.5	65	0.0000
24-hour ventilation requirement	47	79.7	12	20.3	59	0.0000
Respiratory fisioterapia	45	36.6	78	63.4	123	0.0000
Anticoagulant	29	58.0	21	42.0	50	0.0000

Nasogastric tube	57	81.4	13	18.6	70	0.0000
Central Venous Catheter	41	80.4	10	19.6	51	0.0000
Tracheostomy	17	70.8	7	29.2	24	0.0000

Table 3 - Vaccination status and biological risk factors for death due to Influenza A(H1N1)

Risk factor	association measures			
	OR _{crude}	CI 95%	OR _{adjusted}	CI 95%
Biological characteristics				
Female sex	1.417	0.794 - 2.528	1.360	0.718 - 2.591
Age (years)	1.026	1.011 - 1.041	1.020	1.007 - 1.038
Are you obese?	4.091	1.594 - 10.502	3.246	1.221 - 8.632
vaccination status				
Did you vaccinate for Influenza?	1.106	0.345 - 3.540	0.829	0.222 - 3.097
Pre-existing diseases				
Cardiovascular Diseases	3.248	1.266 - 8.333	0.682	0.165 - 2.823
Systemic arterial hypertension	3.636	1.698 - 7.786	2.051	0.769 - 5.470
Cerebrovascular disease	6.760	0.604 - 75.650	1.191	0.056 - 25.050
Congestive heart failure	21.492	2.543 - 181.625	14.425	1.032 - 201.538
Diabetes mellitus	29.538	3.628 - 240.454	10.768	1.075 - 107.819
Respiratory tract diseases	2.301	1.195 - 4.432	1.880	0.855 - 4.134
COPD	3.545	1.200 - 10.468	1.701	0.454 - 6.377
hospital care				
Has it been transferred from service?	3.645	2.101 - 6.324	2.264	1.099 - 4.664
Were you admitted to the ICU?	37.080	17.387 - 79.074	32.742	15.250 - 70.300

Table 4 - Biological, clinical and care risk factors for death due to Influenza A(H1N1)

Risk factor	Model 1		Model 2		Model 3	
	OR _{adjusted}	CI 95%	OR _{adjusted}	CI 95%	OR _{adjusted}	CI 95%
Biological characteristics						
Age (years)	1.015	0.998 - 1.031	1.017	1.000 - 1.035	--	--
Are you obese?	2.631	0.939 - 7.367	2.296	0.785 - 6.708	--	--
Pre-existing diseases						
Congestive heart failure	10.326	1.012 - 105.347	6.851	0.659 - 71.189	--	--
Diabetes mellitus	8.508	0.890 - 81.333	7.122	0.694 - 73.001	24.216	1.772 - 330.811
Respiratory tract diseases	2.182	1.022 - 4.657	2.042	0.929 - 4.490	--	--
hospital care						
Has it been transferred from service?			3.014	1.620 - 5.609	1.897	0.891 - 4.038
Were you admitted to the ICU?					29.468	13.137 - 66.098

Authors' contribution statement

The authors participated in the conception, planning, development, data analysis and writing of the manuscript

Conflict of interest statement

The authors have no affiliation with any organization with a direct or indirect financial interest in the subject matter discussed in the manuscript.

Research data availability statement

The dataset supporting the results of this study is not publicly available.

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.